Please check the examination details below	v before entering your candidate information				
Candidate surname	Other names				
Pearson Edexcel International Advanced Level	re Number Candidate Number				
<b>Monday 20 May</b>	2019				
Morning (Time: 1 hour 30 minutes) Paper Reference <b>WCH11/01</b>					
Chemistry International Advanced Subsidiary/Advanced Level Unit 1: Structure, Bonding and Introduction to Organic Chemistry					
Candidates must have: Scientific cale	<b>Culator</b> Total Marks				

### Instructions

- Use **black** ink or **black** ball-point pen.
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer **all** questions.
- Answer the questions in the spaces provided
  - there may be more space than you need.

### Information

- The total mark for this paper is 80.
- The marks for **each** question are shown in brackets
  - use this as a guide as to how much time to spend on each question.
- You will be assessed on your ability to organise and present information, ideas, descriptions and arguments clearly and logically, including your use of grammar, punctuation and spelling.
- There is a Periodic Table on the back page of this paper.

### **Advice**

- Read each question carefully before you start to answer it.
- Show all your working in calculations and include units where appropriate.
- Check your answers if you have time at the end.

Turn over ▶





## **SECTION A**

# Answer ALL the questions in this section.

You should aim to spend no more than 20 minutes on this section.

For each question, select one answer from A to D and put a cross in the box  $\boxtimes$ . If you change your mind, put a line through the box 🔀 and then mark your new answer with a cross  $\boxtimes$ .

V	۷h	ich	statement is <b>not</b> true for sodium chloride?			
>	■ A sodium chloride conducts electricity in aqueous solution					
>	<	В	sodium chloride conducts electricity when molten			
>	<	C	sodium chloride has a molecular structure			
>	<	D	sodium chloride has a giant structure			
			(Total for Question 1 = 1 mark)			
2 V	۷h	iich	of these molecules is the <b>most</b> polar?			
>	<	A	H—H			
D	<	В	H—F			
Σ	<	C	H—Cl			
Σ	<	D	H—Br			
			(Total for Question 2 = 1 mark)			
3 (	Ξο·	vale	ent bonding is best described as the electrostatic attraction between			
>	<	Α	oppositely charged ions			
>	<	В	positive ions and delocalised electrons			
>	<	c	a shared pair of electrons			
D	<	D	two nuclei and a shared pair of electrons			
			(Total for Question 3 = 1 mark)			
	lca	, +h	is space for any rough working. Anything you write in this space will gain no credit			



**4** Which is correct for tetrafluoromethane (CF<sub>4</sub>)?



	Bonds	Molecule
<b>⋈</b> A	polar	polar
⊠ B	non-polar	polar
⊠ C	polar	non-polar
⊠ D	non-polar	non-polar

(Total for Question 4 = 1 mark)

- 5 Which pair of ions gives the strongest ionic bonding?

  - B K<sup>+</sup> and Br<sup>-</sup>
  - $\blacksquare$  **C** Mg<sup>2+</sup> and O<sup>2-</sup>
  - $\blacksquare$  **D** Ca<sup>2+</sup> and S<sup>2-</sup>

(Total for Question 5 = 1 mark)

6 In 1.31 g of a chloride of titanium, there is 0.528 g of titanium.

What is the empirical formula of this titanium chloride?

[ $A_r$  values: Ti = 47.9 Cl = 35.5]

- A TiCl
- B TiCl₂
- ☑ C TiCl<sub>3</sub>
- ☑ D Ti<sub>2</sub>Cl<sub>4</sub>

(Total for Question 6 = 1 mark)

- 7 Which isotope is used as the standard in the definition of relative atomic mass?
  - $\square$  A  $^{1}$ H


  - C 13C
  - $\square$  **D** <sup>16</sup>O


(Total for Question 7 = 1 mark)


- **8** Which statement about subatomic particles is correct?
  - A neutral atoms always contain the same number of protons and electrons
  - **B** neutral atoms always contain the same number of protons and neutrons
  - $\square$  **C** electrons have a relative mass of 1 and a charge of -1
  - **D** protons have a relative mass of 1 and no charge


(Total for Question 8 = 1 mark)

**9** Which sketch graph shows the trend in first ionisation energy values going down Group 1 in the Periodic Table?










(Total for Question 9 = 1 mark)


- **10** Isotopes are atoms of an element that have different
  - A electronic structures
  - **B** numbers of electrons
  - C numbers of protons
  - **D** numbers of neutrons


(Total for Question 10 = 1 mark)


11 The two stable isotopes of bromine have relative masses of 79 and 81.

Which is the correct pattern of peaks in the mass spectrum of molecular bromine?










(Total for Question 11 = 1 mark)

6



**12** The mass spectrum of a sample of silicon is shown.



What is the **best** estimate for the relative atomic mass of silicon in this sample?

- **■ B** 28.2
- **∠ C** 28.8
- **☑ D** 29.0


(Total for Question 12 = 1 mark)

**13** Which is the equation for the **second** ionisation energy of an element, A?

- $\square$  A A(g)  $\rightarrow$  A<sup>2+</sup>(g) + 2e<sup>-</sup>
- $\blacksquare$  **B**  $A^+(g) \rightarrow A^{2+}(g) + e^-$
- $\square$  **C**  $A^{2+}(g) \rightarrow A^{3+}(g) + e^{-}$
- $\square$  **D**  $A^{2+}(g) \rightarrow A^{4+}(g) + 2e^{-}$

(Total for Question 13 = 1 mark)

**14** The bar chart shows the melting temperatures of the first twenty elements.



The bar chart shows that melting temperatures


- A of giant covalent structures are the highest in Period 2 and in Period 3
- **B** of metals are always higher than non-metals
- C increase going down each group
- D increase across Period 2 and Period 3

(Total for Question 14 = 1 mark)

- **15** Which is the electronic configuration of the Sc<sup>3+</sup> ion?
  - $\triangle$  **A** 1s<sup>2</sup> 2s<sup>2</sup> 2p<sup>6</sup> 3s<sup>2</sup> 3p<sup>6</sup>
  - $\blacksquare$  **B** 1s<sup>2</sup> 2s<sup>2</sup> 2p<sup>6</sup> 3s<sup>2</sup> 3p<sup>5</sup> 3d<sup>1</sup>
  - $\square$  **C** 1s<sup>2</sup> 2s<sup>2</sup> 2p<sup>6</sup> 3s<sup>2</sup> 3p<sup>6</sup> 3d<sup>1</sup> 4s<sup>2</sup>
  - $\square$  **D** 1s<sup>2</sup> 2s<sup>2</sup> 2p<sup>6</sup> 3s<sup>2</sup> 3p<sup>6</sup> 3d<sup>4</sup> 4s<sup>2</sup>

(Total for Question 15 = 1 mark)

16 What is the name of the product when this molecule reacts with chlorine gas?



- A 5,5-dichloro-2-methylpentane
- **B** 4,5-dichloro-2-methylpentane
- ☑ C 2,3-dichloro-4-methylpentane
- ☑ D 1,2-dichloro-4-methylpentane

(Total for Question 16 = 1 mark)

17 What type of bond breaking occurs in this process?

$$H_2O \rightarrow H^+ + OH^-$$

- **A** electrophilic
- ☑ B heterolytic
- C homolytic
- **D** ionic

(Total for Question 17 = 1 mark)

18 What reagent and conditions are used for this conversion?

- A potassium manganate(VII) in aqueous acid
- **B** sodium hydroxide in dilute aqueous solution
- **C** steam and acid catalyst
- D steam and nickel catalyst

(Total for Question 18 = 1 mark)

**19** But-2-ene shows geometric isomerism.

$$H$$
  $H$   $C=C$   $CH_3$ 

What are the prefixes that could be used in naming this isomer?

	cis / trans	E/Z
⊠ A	cis	Ε
<b>⋈</b> B	cis	Z
⊠ C	trans	Е
⊠ D	trans	Z

(Total for Question 19 = 1 mark)

**20** Which is the major product of this reaction?

$$H \longrightarrow Br$$
  $+$   $C \longrightarrow C$   $\rightarrow$   $H$ 

(Total for Question 20 = 1 mark)

**TOTAL FOR SECTION A = 20 MARKS** 

### **SECTION B**


# Answer ALL the questions. Write your answers in the spaces provided.

- 21 Magnesium carbonate powder reacts with hydrochloric acid.
  - (a) Complete the equation for this reaction by adding state symbols.

(1)

$$MgCO_3(.....) + 2HCl(....) \rightarrow MgCl_2(....) + H_2O(....) + CO_2(....)$$

(b) A student carried out an investigation to determine the molar volume of carbon dioxide using this apparatus.




The student carried out five experiments, adding a different mass of magnesium carbonate each time.


The results are shown in the table.

Mass of magnesium carbonate / g	Volume of gas collected / cm <sup>3</sup>
0.05	11
0.10	27
0.15	38
0.20	54
0.25	63

(i) Plot a graph of these results.



(ii) A student carried out a further experiment using a different mass of magnesium carbonate.



Give the volume of gas collected using the **inverted** measuring cylinder.

(1)

(3)



(iii) Determine the mass of magnesium carbonate added in the experiment in (b)(ii), using your graph.

(1)

(iv) Calculate the molar volume of carbon dioxide using your answers to parts (b)(ii) and (b)(iii). Give your value to an appropriate number of significant figures and include units.

[ $A_r$  values: Mg = 24.3 C = 12.0 O = 16.0]

(4)

(v) The acid must be in excess for each experiment.

Calculate the **minimum** concentration of hydrochloric acid needed for 30 cm<sup>3</sup> of acid to completely react with 0.25 g of magnesium carbonate.

$$MgCO_3 + 2HCl + MgCl_2 + H_2O + CO_2$$
(2)

(c) The value of molar volume calculated in (b)(iv) was lower than the student expected.

Give **two** reasons for the value being lower than expected.

Assume that the correct amounts of hydrochloric acid and magnesium carbonate were used.

(2)

(Total for Question 21 = 14 marks)



22	This a	westion is about fivele and not more				
22	2 This question is about fuels and polymers.					
	Used coffee grounds have been suggested as a carbon-neutral fuel to replace some fossil fuels.					
	(a) (i)	Explain why coffee grounds might be considered a carbon-neutral fuel.	(2)			
	(ii)	Explain how the use of fossil fuels causes climate change.	(2)			

<ul><li>(b) Long chain alkanes are not normally used as fuels as they produce soot.</li><li>(i) Name another pollutant formed by <b>incomplete</b> combustion of alkanes.</li></ul>	(1)
(ii) Write the equation for the <b>complete</b> combustion of octane. State symbols are not required.	(2)
<ul><li>(c) Long chain alkanes are converted into smaller, more useful molecules including</li><li>(i) Name this process.</li></ul>	alkenes.
(ii) Give a test for alkenes, including the positive result.	(2)
(d) Alkenes, such as ethene, can be used to make polymers.  (i) Write a balanced equation for the polymerisation of ethene using displayed formulae.	(1)



(ii) Bananas produce ethene as they ripen.	
Suggest one advantage and one disadvantage of using ripening bananas a source of ethene for polymer production.	is a
source of earlier for polymer production.	(2)
<ul><li>(e) Burning poly(chloroethene) in an incinerator results in the formation of hydrogen chloride.</li></ul>	
(i) State a <b>hazard</b> associated with hydrogen chloride.	(1)
	(1)
(ii) Suggest how the hydrogen chloride could be removed from the waste gase	PS
produced in an incinerator.	(1)
	(-)
(Total for Question 22 = 15	marks)

- 23 This question is about bonding.
  - (a) Draw an electron density map for a molecule of oxygen.

(1)

(b) Draw a diagram to show the shape of a water molecule. Give the bond angle.


(2)

Bond angle .....

- (c) The compound POCl₃ has a simple molecular structure.
  - (i) Complete the dot-and-cross diagram for the POCl<sub>3</sub> molecule.

Use crosses (x) for the phosphorus electrons, dots (●) for the chlorine electrons and circles (o) for the oxygen electrons.

(2)




(ii) Explain the shape of this molecule using the electron-pair repulsion theory.

(3)

١
)

- (e) Diamond, graphite and graphene are all forms of carbon.
  - (i) Explain **two** ways in which the physical properties of diamond and graphite differ. Refer to their structure and bonding in your answer.

(4)





graphite


(ii) State how the structure of graphene is related to the structure of graphite. (1)





(Total for Question 23 = 20 mar	
(iii) State a use for graphene, identifying the property that makes it suitable for that	use. (2)

**24** Airbags protect occupants by inflating when a car crashes.

Airbags rely on chemical reactions to produce large volumes of gases quickly. In some airbags, solid sodium azide (NaN<sub>3</sub>) decomposes forming nitrogen gas and sodium as the only products.

(a) Write an equation for the decomposition of sodium azide. State symbols are not required.

(1)

(b) A passenger airbag requires 120 dm<sup>3</sup> of gas to fill it.

Calculate, using the ideal gas equation, the mass of sodium azide required to fill a passenger airbag in this reaction under standard conditions (101 000 Pa, 25°C).

Give your answer to an appropriate number of significant figures.

$$[pV = nRT$$
  $R = 8.31 \,\mathrm{J}\,\mathrm{K}^{-1}\,\mathrm{mol}^{-1}]$ 

(6)

(c) Two further reactions take place in the airbag.

Reaction **A** 10Na + 2KNO<sub>3</sub> 
$$\rightarrow$$
 K<sub>2</sub>O + 5Na<sub>2</sub>O + N<sub>2</sub>

Reaction **B** 
$$K_2O + Na_2O + SiO_2 \rightarrow Na_2K_2SiO_4$$

(i) Reaction A produces more nitrogen to inflate the airbag.

Calculate the atom economy, by mass, for the production of nitrogen in reaction **A**. Give your answer to an appropriate number of significant figures.

(3)

(ii) State the type of reaction taking place in reaction **B**.

(1)

(Total for Question 24 = 11 marks)

TOTAL FOR SECTION B = 60 MARKS TOTAL FOR PAPER = 80 MARKS



# **BLANK PAGE**

# **BLANK PAGE**



# The Periodic Table of Elements

0 (8)	(18) 4.0 <b>He</b> helium 2	
7	(71)	
9	(16)	
2	(15)	,
4	(13) (14)	,
m	(13)	
	1.0  H hydrogen 1	
2	(2)	
_	$\mathcal{E}$	` `

1	20.2	Ne	neon	10	39.9	Αľ	argon	9	83.8	궃	krypton	36	131.3	Xe	xenon	54	[222]	R	radon	98		ted		
(,,,)	19.0	ш	fluorine	6	35.5	บ	chlorine	17	79.9	Br	bromine	35	126.9	Ι	iodine	53	[210]	At	astatine	85		een repor		
(0.1)	16.0	0	oxygen	∞	32.1	S	sulfur	16	79.0	Se	selenium	34	127.6	<u>P</u>	tellurium	52	[506]	S	polonium	84		116 have b	ticated	
(61)	14.0	z	nitrogen	7	31.0	۵	phosphorus	15	74.9	As			121.8			51	209.0	B:	bismuth	83		nbers 112-	but not fully authenticated	
(, , )	12.0	U	carbon	9	28.1			4	72.6	ge	germanium	32	118.7	Sn			207.2	Ъ	lead	82		atomic nur	but not fi	
(5.)	10.8	മ	boron	2	27.0	¥	aluminium	13	2.69		gallium	31	114.8	I	indium	49	204.4	F	thallium	81		Elements with atomic numbers 112-116 have been reported		
								(71)	65.4	Zn	zinc	30	112.4	<u>გ</u>	cadminm	48	200.6	Ξğ	mercury	80		Elem		
							( )	(1.1)	63.5	D C	copper	29	107.9	Ag	silver	47	197.0	Αu	plog	79	[272]	Rg	roentgenium	111
							3	(01)	58.7	'n	nickel	28	106.4	Pq	palladium	46	195.1	꿉	platinum	78	[271]	Ds	darmstadtium	110
							ę	(%)	58.9	ပိ	cobalt	27	102.9	R	rhodium	45	192.2	<u>_</u>	iridium	77	[568]	Mt	Щ	109
							Ć	(&)	55.8	Fe	iron	56	101.1	Ru	ruthenium	44	190.2	o	osmium	76	[277]	Hs		108
•							į	$\leq$	54.9	Wn	manganese	25	[86]	ပ	technetium	43	186.2	Re	rhenium	75	[564]	Bh	bohrium	107
	mass	lod		nmber			S	(0)	52.0	င်	chromium	24 25	95.9	Wo	molybdenum technetium	42	183.8	>	tungsten	74	[596]	Sg	seaborgium	106
	relative atomic mass	atomic symbol	name	atomic (proton) number			į	(c)	50.9	>	vanadium	23	92.9	g	niobium	41	180.9	٦a	tantalum	73	[797]		Ε	105
	relati	ato		atomic			5	(4)	47.9	ï	titanium	22	91.2	Zr	zirconium	40	178.5	Ŧ	hafnium	72	[261]	Ř	rutherfordium	104
							Ć	(5)	45.0	Sc	scandium	21	6.88	>	yttrium	39	138.9	La*	lanthanum	22	[227]	Ac*	actinium	88
(-)	0.6	Be	beryllium	4	24.3	Mg	magnesium	12	40.1	Ca	calcinm	20	9.78	Sr	strontium	38	137.3	Ba	barium	99	[526]	Ra	radium	88
	6.9	<u></u>	lithium	m	23.0	Na	sodium	1	39.1	¥	potassium	19	85.5	&	rubidium	37	132.9	S	caesium	22	[223]	Ŧ	francium	87

\* Lanthanide series

\* Actinide series

			_			_	
175	Γn	lutetium	71	[257]	۲	lawrencium	103
173	ХÞ	ytterbium	70	[254]	8	nobelium	102
169	Ε	thulium	69	[526]	Þ₩	mendelevium	101
167	ם	erbium	89	[253]	Fn	fermium	100
165	운	holmium	29	[254]	Es	einsteinium	66
163	کم	dysprosium	99	[251]	ರ	californium	98
159	ТР	terbium	65	[245]	쓞	perkelium	26
157	В	gadolinium	64	[247]	CH	aurium	96
152	Eu	europium	63	[243]	Am	americium	95
150	Sm	samarinm	62	[242]	Pu	plutonium	94
[147]	Pm	promethium	61	[237]	å	neptunium	93
144	PN	neodymium	90	238	_	uranium	92
141	P			[231]	Ра	protactinium	91
140	Ce	cerium	58	232	Ļ	thorium	90