



# Mark Scheme (Results)

October 2023

Pearson Edexcel International Advanced  
Subsidiary Level In Chemistry (WCH13)  
Paper 01  
Unit 3: Practical Skills in Chemistry I

## **Edexcel and BTEC Qualifications**

Edexcel and BTEC qualifications are awarded by Pearson, the UK's largest awarding body. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information visit our qualifications websites at [www.edexcel.com](http://www.edexcel.com) or [www.btec.co.uk](http://www.btec.co.uk). Alternatively, you can get in touch with us using the details on our contact us page at [www.edexcel.com/contactus](http://www.edexcel.com/contactus).

## **Pearson: helping people progress, everywhere**

Pearson aspires to be the world's leading learning company. Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: [www.pearson.com/uk](http://www.pearson.com/uk)

October 2023

Question Paper Log Number: P75062A

Publications Code: WCH13\_01\_MS\_2310

All the material in this publication is copyright.

© Pearson Education Ltd 2023

## General Marking Guidance

- All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.
- Mark schemes should be applied positively. Candidates must be rewarded for what they have shown they can do rather than penalised for omissions.
- Examiners should mark according to the mark scheme not according to their perception of where the grade boundaries may lie.
- There is no ceiling on achievement. All marks on the mark scheme should be used appropriately.
- All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e. if the answer matches the mark scheme. Examiners should also be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.
- Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.
- When examiners are in doubt regarding the application of the mark scheme to a candidate's response, the team leader must be consulted.
- Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.


| Question Number | Answer  | Additional Guidance  | Mark |
|-----------------|---|--|------|
| 1(a)(i)         | <p>A description that makes reference to the following points:</p> <ul style="list-style-type: none"> <li>• use of nichrome / platinum / Pt wire</li> <li>• use of (concentrated) hydrochloric acid / HCl(aq)</li> <li>• flame test method</li> </ul> | <p>(1) Allow nickel-chromium / Ni-Cr<br/>Allow silica rod<br/>Allow rod for wire<br/>Do not award just 'nickel' / just 'chromium'<br/>Do not award wooden splint</p> <p>(1) Allow any mention of HCl(aq) e.g. cleaning or mixing solid and acid or making a paste / solution<br/>Allow HCl for HCl(aq)<br/>Ignore dilute<br/>Do not award other acids</p> <p>(1) (wire then) dipped in solid<br/><b>and</b><br/>(placed) in (hot / roaring / colourless / blue-cone / non-luminous) (Bunsen) <b>flame</b></p> <p>Allow salt / compound / substance / paste / sample / solution for 'solid'<br/>Allow on / over / under / near / show / above for 'in'<br/>Allow spirit / ethanol burner</p> <p>Do not award 'metal' for solid<br/>Do not award fire for flame<br/>Do not award yellow / luminous flame<br/>Ignore 'burn'</p> | (3)  |

| Question Number | Answer  | Additional Guidance   | Mark |
|-----------------|---|---|------|
| 1(a)(ii)        | <p>An answer that makes reference to the following points:</p> <ul style="list-style-type: none"> <li>statement of both flame colours</li> <li>identification of Ba<sup>2+</sup></li> </ul> | <p>(1) Na<sup>+</sup> = (persistent) yellow<br/> Allow gold / orange / yellow-orange<br/> K<sup>+</sup> = lilac<br/> Allow (pale) purple<br/> Do not award violet</p> <p>(1) Ignore barium / barium ion<br/> Do not award Cu<sup>2+</sup></p> | (2)  |

| Question Number | Answer  | Additional Guidance   | Mark |
|-----------------|---|---|------|
| 1(b)(i)         | <p>An answer that makes reference to the following:</p> <ul style="list-style-type: none"> <li>chloride / Cl<sup>-</sup> <b>and</b> bromide / Br<sup>-</sup> <b>and</b> iodide / I<sup>-</sup></li> </ul> | <p>Allow omission of the charge once<br/> Ignore chlorine ion, bromine ion or iodine ion<br/> Do not award just chlorine, bromine or iodine</p> | (1)  |

| Question Number | Answer  | Additional Guidance   | Mark |
|-----------------|---|---|------|
| 1(b)(ii)        | <p>An answer that makes reference to the following points:</p> <p>EITHER</p> <ul style="list-style-type: none"> <li>• addition of <b>dilute</b> (aqueous) ammonia (1)</li> <li>• only silver chloride / chloride (precipitate) dissolves (1)</li> <li>• addition of <b>concentrated</b> (aqueous) ammonia (1)</li> <li>• only silver iodide / iodide is insoluble (1)</li> </ul> <p>OR</p> <ul style="list-style-type: none"> <li>• addition of <b>concentrated</b> sulfuric acid /H<sub>2</sub>SO<sub>4</sub> (1)</li> <li>• silver chloride / chloride precipitate gives steamy / misty / white fumes only (1)</li> <li>• silver bromide / bromide precipitate gives brown fumes (1)</li> <li>• silver iodide / iodide precipitate gives purple vapour fumes / gas (1)</li> </ul> | <p>Penalise use of halogen for halide once only<br/>Allow use of halogen ion / anion<br/>The sequence suggested must allow the ions to be distinguished</p> <p>Allow bromide and iodide do not dissolve</p> <p>Accept only silver bromide / bromide precipitate dissolves if chloride has been eliminated and conc ammonia added to remaining precipitates</p> <p>If no other mark is scored ‘addition of (aqueous) ammonia scores (1)</p> <p>Allow addition to A, B and C for this set of tests</p> <p>Allow choking gas evolved / choking smell</p> <p>Allow black / grey solid or ‘bad eggs’ smell<br/>Allow yellow solid (of sulfur)<br/>Ignore choking gas evolved / choking smell</p> <p>If no other mark is scored, silver chloride is white, <b>and</b> silver bromide is cream <b>and</b> silver iodide is yellow scores (1)</p> | (4)  |

(Total for Question 1 = 10 marks)

| Question Number | Answer   | Additional Guidance  | Mark |
|-----------------|--|--|------|
| 2(a)            | <p>Diagram showing the following points</p> <ul style="list-style-type: none"> <li>• boiling tube</li> <li>• delivery tube above the level of the liquid in the boiling tube if shown or above half the height of the tube</li> <li>• sealed connection between reaction vessel and delivery tube</li> <li>• inverted measuring cylinder containing some water (shown as a line or an annotation indicating it is full)</li> <li>• water level in trough over the bottom of the measuring cylinder</li> <li>• end of delivery tube below or in the measuring cylinder</li> </ul> <p>6 points scores 3 marks;<br/>5 or 4 points scores 2 marks;<br/>2 or 3 points scores 1 mark</p> | <p>Allow any reaction apparatus that can be fitted with a bung and delivery tube<br/>Allow any part of a delivery tube shown</p> <p>Allow bung not shown as cross-section</p> <p>Allow measuring cylinder shown not vertical<br/>Allow other collecting vessels with graduations</p> <p>Ignore omission of graduations on measuring cylinder<br/>Ignore omission of beehive shelf<br/>Ignore gas syringe</p> <p>Ignore the delivery tube passing through the wall of the trough</p> <p style="text-align: center;">Alternative allowed bungs</p>  <p>The diagram shows a boiling tube containing blue liquid, with a delivery tube extending from its mouth into an inverted measuring cylinder. The measuring cylinder is partially filled with blue liquid and is submerged in a trough of water. To the right, two alternative bung designs are shown: a solid orange square and a square with a vertical slot.</p> <p>Ignore apparatus labels including 'heat'</p> | (3)  |

| Question Number | Answer   | Additional Guidance   | Mark |
|-----------------|--|---|------|
| 2(b)            | <p>An explanation that makes reference to the following points:</p> <ul style="list-style-type: none"> <li>ethanoic acid is a weak acid (and hydrochloric acid is a strong acid) (1)</li> <li>(with hydrochloric acid)</li> </ul> <p>gas would escape before the boiling tube was sealed (1)</p> | <p>Allow 'hydrochloric acid is a <b>stronger</b> acid'</p> <p>Allow reaction would be slow</p> <p>Allow reaction would be less vigorous</p> <p>Allow reaction would not be violent</p> <p>Allow reverse arguments</p> <p>Ignore just 'hydrochloric acid is a strong acid'</p> <p>Allow just 'less gas would escape'</p> | (2)  |

| Question Number | Answer   | Additional Guidance   | Mark |
|-----------------|--|---|------|
| 2(c)            | <p>An answer that makes reference to the following point:</p> <ul style="list-style-type: none"> <li>to take account of any calcium carbonate left in the weighing bottle</li> </ul> | <p>Ignore just 'gives the mass of CaCO<sub>3</sub> that reacts'</p> <p>Ignore just 'gives exact amount of CaCO<sub>3</sub>'</p> | (1)  |

| Question Number | Answer   | Additional Guidance  | Mark  |
|-----------------|--|--|---|
| 2(d)(i)         | <ul style="list-style-type: none"> <li>• correct axes with at least 50% of the grid used in both directions (1)</li> <li>• axes labelled with correct units (1)</li> <li>• all points plotted correctly (1)</li> </ul> | <p>mass on x axis; volume on y axis<br/>Allow scale 0.10 g = 1 big square</p> <p>Allow grammatical errors e.g. use of brackets rather than ‘/’ before units<br/>Allow mass / g and vol / cm<sup>3</sup> for labels</p> <p>Allow plotting to within half a small square</p> <p>TE for M2 and M3 if axes wrong way round</p> | <p>(3)</p> <p><b>Clip with d(ii) and (d)(iii)</b></p> |

| Question Number | Answer  | Additional Guidance   | Mark   |
|-----------------|---|---|--|
| 2(d)(ii)        | <p>An answer that makes reference to the following points:</p> <ul style="list-style-type: none"> <li>• best fit line drawn through five points <b>and</b> passes through the origin</li> <li>• structure line from 0.25 g to graph <b>and</b> line to 52.5 cm<sup>3</sup></li> </ul> | <p>Do not award BFL drawn to include value at 0.2 g<br/> Allow TE on points plotted in (d)(i) even if it does not go through origin<br/> Accept line that stops at 0.11 g data point but would pass through the origin if extrapolated<br/> Ignore extrapolation beyond 0.36 / 76</p> <p>Allow vol at 0.25 g marked on BFL<br/> Allow 51.5-53.5 cm<sup>3</sup><br/> TE only on any <b>straight</b> line covering <b>all</b> the points</p> <p>Ignore <b>calculation</b> of volume</p> | <p>(2)</p> <p><b>Clip with d(i) and d(iii)</b></p> |

| Question Number | Answer  | Additional Guidance   | Mark  |
|-----------------|---|---|---|
| 2(d)(iii)       | <ul style="list-style-type: none"> <li>• calculation of molar mass of CaCO<sub>3</sub> (1)</li> <li>• calculation of amount of CaCO<sub>3</sub> (1)</li> <li>• scale volume of CO<sub>2</sub> to 1 mol (1)</li> </ul> | <p>Example of calculation</p> <p>molar mass = <math>40.1 + 12 + 16 \times 3 = 100.1 \text{ (g mol}^{-1}\text{)}</math></p> <p>mol (CaCO<sub>3</sub>) = <math>0.25 \div 100.1 = 2.4975 \times 10^{-3} / 0.0024975</math></p> <p>1 mol CO<sub>2</sub> occupies <math>52.5 \div 2.4975 \times 10^{-3} = 21021 \text{ (cm}^3\text{)} / 21.021 \text{ dm}^3</math><br/> Units must be correct if given but accept <math>\text{cm}^3 \text{ mol}^{-1} / \text{dm}^3 \text{ mol}^{-1}</math></p> <p>TE at each stage and on volume in (d)(ii)</p> <p>Ignore SF except 1 SF</p> <p>Correct answer with some working scores (3)</p> <p>Allow use of <math>M_r</math> (CaCO<sub>3</sub>) = 100 (<math>V_m = 21</math>)</p> <p>Use of 51.5 gives 20621 cm<sup>3</sup> (20600 for <math>M_r = 100</math>)<br/> Use of 52 gives 20821 cm<sup>3</sup> (20800 for <math>M_r = 100</math>)<br/> Use of 53 gives 21221 cm<sup>3</sup> (21200 for <math>M_r = 100</math>)<br/> Use of 53.5 gives 21421 cm<sup>3</sup> (21400 for <math>M_r = 100</math>)</p> <p>Calculation of moles of acid (0.030) divided by 2 (0.015)<br/> Vol of CO<sub>2</sub> <math>\div 0.015</math> e.g. <math>52.5 \div 0.015 = 3500 \text{ cm}^3</math> scores M3 only</p> | <p>(3)</p> <p><b>Clip with<br/>d(i) and<br/>d(ii)</b></p> |



| Question Number | Answer  | Additional Guidance   | Mark  |
|-----------------|---|---|---|
| 3(a)(i)         | <p>An answer that makes reference to the following points:</p> <ul style="list-style-type: none"> <li>flammable symbol identified</li> </ul> <p><b>and</b></p> <ul style="list-style-type: none"> <li>harmful to the environment symbol identified</li> </ul> | <p>Allow inflammable<br/>Ignore combusts / burns easily</p> <p>Allow alternatives to 'harmful to' e.g. 'damages' / 'bad for' / 'poisonous' / 'toxic' / 'hazard' to the environment<br/>Allow 'living things' / 'organisms' for 'the environment'<br/>Ignore type of environment e.g. aquatic<br/>Ignore pollutes the environment</p> <p>Do not award 'biohazard'<br/>Do not award symbols the wrong way round<br/>Do not award just 'toxic' / 'poisonous' / 'harmful'</p> | <p><b>(1)</b></p> <p><b>Clip with (a)(ii)</b></p> |

| Question Number | Answer   | Additional Guidance   | Mark              |
|-----------------|--|---|-------------------|
| 3(a)(ii)        | <p>An answer that makes reference to the following points:</p> <ul style="list-style-type: none"> <li>(cyclohexanol(flammable)) use an electric heater <b>(1)</b></li> <li>(cyclohexene (harmful to the environment)) use an organic waste bottle / separate container <b>(1)</b></li> </ul> | <p>Ignore use of fume cupboard, goggles, lab coat</p> <p>Accept isomantle<br/>Allow water bath / oil bath<br/>Allow 'no (naked) flame / fire'<br/>Ignore 'keep away from oxidising agents'<br/>Ignore keep away from Bunsen burner</p> <p>Accept do <b>not</b> pour the (organic) waste down the sink / drain<br/>Allow do not release into the environment</p> <p>Allow one use of 'use small amounts' in M1 or M2<br/>No TE on incorrect identification of hazard symbols</p> | <p><b>(2)</b></p> |

| Question Number | Answer   | Additional Guidance  | Mark |
|-----------------|--|--|------|
| 3(b)            | <p>An answer that makes reference to the following:</p> <ul style="list-style-type: none"> <li>(anti-bumping granules) provides a nucleus on which gas bubbles grow</li> </ul> | <p>Accept provide a surface /site for bubbles to form on</p> <p>Allow prevent local heating / superheating</p> <p>Allow 'distribute the heat'</p> <p>Allow prevent the (sudden) production of large gas bubbles (which cause bumping)</p> <p>Allow ensures that gas bubbles are small</p> <p>Ignore stir the reaction mixture</p> <p>Ignore prevent flash boiling / sudden boiling</p> | (1)  |

| Question Number | Answer   | Additional Guidance   | Mark |
|-----------------|--|---|------|
| 3(c)            | <p>An answer that makes reference to the following points:</p> <ul style="list-style-type: none"> <li>(fractional distillation) gives better separation of the substances in the reaction flask (than simple distillation)</li> <li>(fractional distillation column) gives longer reaction time</li> </ul> <p>OR</p> <ul style="list-style-type: none"> <li>'better separation of cyclohexene and water' scores 2 marks</li> </ul> | <p>(1) Allow (fractional distillation) is more effective / efficient (than simple distillation)</p> <p>Allow just 'better separation' is achieved</p> <p>Allow purer product obtained</p> <p>Ignore reference to increased yield</p> <p>Ignore 'more accurate'</p> <p>Ignore 'similar boiling temperatures'</p> <p>(1) Accept fractional distillation column acts like a reflux condenser</p> <p>Accept cyclohexanol has a higher boiling temperature (than cyclohexene) so is returned to the flask, increasing reaction time / yield</p> <p>Allow to give (more) complete reaction</p> <p>Just 'separation of cyclohexene and water' scores (1)</p> | (2)  |



| Question Number | Answer   | Additional Guidance   | Mark |
|-----------------|--|---|------|
| 3(e)            | <p>An explanation that makes reference to the following points:</p> <ul style="list-style-type: none"> <li>(the organic layer) changes from cloudy to clear</li> <li>(anhydrous calcium chloride) removes the (traces of) water</li> </ul> | <p>(1) Allow just 'becomes clear / transparent'<br/>Ignore 'less cloudy', clearer, more transparent<br/>Ignore white<br/>Ignore colourless<br/>Ignore the drying agent clumps together<br/>Ignore layers disappear</p> <p>(1) Accept (anhydrous calcium chloride) dries the cyclohexene<br/>Allow (anhydrous) calcium chloride is a drying agent<br/>Ignore calcium chloride becomes hydrated</p> | (2)  |

| Question Number | Answer   | Additional Guidance   | Mark |
|-----------------|--|---|------|
| 3(f)            | <p>An answer that makes reference to the following point:</p> <ul style="list-style-type: none"> <li>lower temperature within a suitable range<br/><b>and</b><br/>upper temperature within a suitable range</li> </ul> | <p>79–82 (°C)</p> <p>84–88 (°C)</p> <p>Do not award a range which starts or ends with 83 (°C)</p> | (1)  |

| Question Number | Answer   | Additional Guidance  | Mark |
|-----------------|--|--|------|
| 3 (g)           | <ul style="list-style-type: none"> <li>• calculation of mass of cyclohexene formed from 3.96 g of cyclohexanol with 100% yield (1)</li> <li>• calculation of percentage yield (1)</li> </ul> | <p>Example of calculation</p> <p>100 g of cyclohexanol forms 82 g of cyclohexene and<br/>(so) 3.96 g forms <math>3.96 \times 82 \div 100 = 3.2472</math> (g)</p> <p>% yield = <math>100 \times 2.09 \div 3.2472 = 64.363\%</math></p> <p>Method using calculation of moles<br/>mol of cyclohexanol = <math>3.96 \div 100</math><br/>= <math>3.96 \times 10^{-2} / 0.0396</math><br/>= mol cyclohexene<br/>mass of cyclohexene for 100% yield<br/>= <math>82 \times 0.0396 = 3.2472</math> (g) (1)<br/>% yield = <math>100 \times 2.09 \div 3.2472</math><br/>= <math>64.363\%</math> (1)</p> <p>Ignore SF except 1 SF</p> <p>TE for <b>numerical</b> errors in M1 unless %&gt;100</p> <p>Allow any correct method<br/>Correct answer with some working scores (2)</p> <p><math>100 \times 2.09 \div 3.96 = 52.778\%</math> scores zero</p> | (2)  |

| Question Number                    | Answer  | Additional Guidance  | Mark |              |  |              |             |                                    |                              |  |                           |  |   |     |
|------------------------------------|---|--|------|--------------|--|--------------|-------------|------------------------------------|------------------------------|--|---------------------------|--|---|-----|
| 3(h)                               | <p>An answer that makes reference to the following points:</p> <ul style="list-style-type: none"> <li>any three correct observations (1)</li> <li>fourth correct observation (1)</li> </ul> | <table border="1" data-bbox="1189 304 1924 799"> <thead> <tr> <th data-bbox="1189 304 1413 373" rowspan="2">Test</th> <th colspan="2" data-bbox="1413 304 1924 373">Observations</th> </tr> <tr> <th data-bbox="1413 373 1666 442">cyclohexanol</th> <th data-bbox="1666 373 1924 442">cyclohexene</th> </tr> </thead> <tbody> <tr> <td data-bbox="1189 442 1413 620">addition of phosphorus(V) chloride</td> <td data-bbox="1413 442 1666 620">steamy / misty / white fumes</td> <td data-bbox="1666 442 1924 620">no change / no reaction / no observation</td> </tr> <tr> <td data-bbox="1189 620 1413 799">addition of bromine water</td> <td data-bbox="1413 620 1666 799">brown / orange / yellow Br<sub>2</sub>(aq) unchanged</td> <td data-bbox="1666 620 1924 799">brown / orange / yellow Br<sub>2</sub>(aq) turns colourless</td> </tr> </tbody> </table> <p data-bbox="1178 839 1637 871">For PCl<sub>5</sub> do not award white smoke</p> <p data-bbox="1178 916 1834 979">For Br<sub>2</sub>(aq) and cyclohexanol allow no change / no reaction / no observation</p> <p data-bbox="1178 1024 1877 1088">For Br<sub>2</sub>(aq) and cyclohexene allow just 'decolorised' / 'turns colourless'</p> <p data-bbox="1178 1101 1845 1133">Do not award red or red-brown for colour of Br<sub>2</sub>(aq)</p> <p data-bbox="1178 1171 1733 1203">Do not award additional incorrect observations</p> | Test | Observations |  | cyclohexanol | cyclohexene | addition of phosphorus(V) chloride | steamy / misty / white fumes | no change / no reaction / no observation | addition of bromine water | brown / orange / yellow Br <sub>2</sub> (aq) unchanged | brown / orange / yellow Br <sub>2</sub> (aq) turns colourless | (2) |
| Test                               | Observations  |  |      |              |  |              |             |                                    |                              |  |                           |  |   |     |
|                                    | cyclohexanol  | cyclohexene  |      |              |  |              |             |                                    |                              |  |                           |  |   |     |
| addition of phosphorus(V) chloride | steamy / misty / white fumes  | no change / no reaction / no observation   |      |              |  |              |             |                                    |                              |  |                           |  |   |     |
| addition of bromine water          | brown / orange / yellow Br <sub>2</sub> (aq) unchanged  | brown / orange / yellow Br <sub>2</sub> (aq) turns colourless  |      |              |  |              |             |                                    |                              |  |                           |  |   |     |

(Total for Question 3 = 15 marks)

| Question Number | Answer   | Additional Guidance   | Mark |
|-----------------|--|---|------|
| 4(a)            | An answer that makes reference to the following points: <ul style="list-style-type: none"> <li data-bbox="371 347 1207 384">• (from) yellow (1)</li> <li data-bbox="371 421 1207 458">• (to) orange (1)</li> </ul> | Do not award 'red / pink'<br><br>From orange to yellow scores (1) | (2)  |

| Question Number | Answer   | Additional Guidance           | Mark |
|-----------------|--|-------------------------------|------|
| 4(b)            | An answer that makes reference to the following: <ul style="list-style-type: none"> <li data-bbox="371 756 1207 793">• 23.40 (cm<sup>3</sup>)</li> </ul> | Allow 23.4 (cm <sup>3</sup> ) | (1)  |

| Question Number | Answer   | Additional Guidance   | Mark                                   |
|-----------------|--|---|--|
| 4(c)            | <p>A description that makes reference to the following points:</p> <ul style="list-style-type: none"> <li>• add the acid (quickly) until just short of the rough end-point</li> <li>• add the acid drop-by-drop</li> <li>• with swirling</li> </ul> <p><b>and</b></p> <p>until the indicator colour (just) changes</p> | <p>(1) Accept within 1– 4 cm<sup>3</sup> (any value in this range) before the rough titre<br/> Allow to a value in the range 19.4 – 22.4 cm<sup>3</sup><br/> Allow ‘until close to the rough value’<br/> Ignore ‘carbonate added’<br/> Do not award ‘until the rough value is reached’</p> <p>(1) Standalone mark (award even if M1 not given)<br/> Allow ‘dropwise’<br/> Allow ‘a few drops at a time’<br/> Ignore ‘add very slowly’</p> <p>Allow any indication of mixing<br/> Allow shaking<br/> Allow stirring</p> <p>(1) Allow until the end-point<br/> Allow any stated final colour</p> <p>Ignore references to filling the burette, use of the pipette, white tiles</p> | <p>(3)</p> <p><b>Clip with (b)</b></p> |

| Question Number | Answer  | Additional Guidance  | Mark |
|-----------------|---|--|------|
| 4(d)            | <ul style="list-style-type: none"> <li>• calculation of moles of sodium carbonate (1)</li> <li>• use of 2:1 ratio to gives moles of HCl in 22.65 cm<sup>3</sup> (1)</li> <li>• concentration of hydrochloric acid in mol dm<sup>-3</sup> (1)</li> </ul> | <p>Example of calculation</p> $\text{mol Na}_2\text{CO}_3 = 25.0 \times 0.105 \div 1000$ $= 2.625 \times 10^{-3} / 0.002625$ $2 \times 2.625 \times 10^{-3} = 5.25 \times 10^{-3}$ $5.25 \times 10^{-3} \times 1000 \div 22.65$ $= 0.23179 / 2.3179 \times 10^{-1} \text{ (mol dm}^{-3}\text{)}$ <p>The reacting volumes transposed gives the final concentration of hydrochloric acid = 0.19026 (mol dm<sup>-3</sup>). This scores (2)</p> <p>The same volume used twice will give the final concentration of hydrochloric acid = 0.2100 (mol dm<sup>-3</sup>). These score (2)</p> <p>TE at each stage<br/>Ignore SF except 1 SF<br/>Allow any correct method<br/>Correct answer with some working scores (3)</p> <p>Comment<br/>If working shows factor of 1000 omitted twice in the step by step calculation max (2)</p> | (3)  |

**(Total for Question 4 = 9 marks)**  
**(Total for Paper = 50 marks)**

