Please check the examination details b Candidate surname	elow before ent	Other names
Pearson Edexcel International Advanced Level	entre Number	Candidate Number
Wednesday 16	Janu	ary 2019
Morning (Time: 1 hour 30 minutes)	Paper F	Reference WCH02/01
Chemistry Advanced Subsidiary Unit 2: Application of Cor	e Princip	les of Chemistry
Candidates must have: Scientific o	calculator	Total Marks

Instructions

- Use **black** ink or **black** ball-point pen.
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer **all** questions.
- Answer the questions in the spaces provided
 - there may be more space than you need.

Information

- The total mark for this paper is 80.
- The marks for **each** question are shown in brackets
 - use this as a guide as to how much time to spend on each question.
- Questions labelled with an asterisk (*) are ones where the quality of your written communication will be assessed
 - you should take particular care with your spelling, punctuation and grammar, as well as the clarity of expression, on these questions.
- A Periodic Table is printed on the back cover of this paper.

Advice

- Read each question carefully before you start to answer it.
- Show all your working in calculations and include units where appropriate.
- Check your answers if you have time at the end.

Turn over ▶



P 5 4 3 4 2 A 0 1 2 4

SECTION A

Answer ALL the questions in this section. You should aim to spend no more than 20 minutes on this section. For each question, select one answer from A to D and put a cross in the box ⊠. If you change your mind, put a line through the box ⋈ and then mark your new answer with a cross ⋈.

- 1 Which of these molecules is planar?
 - A Ethane, CH₃CH₃
 - ☑ B Ethanoic acid, CH₃COOH
 - C Methanal, HCHO
 - ☑ D Methanol, CH₃OH

(Total for Question 1 = 1 mark)

- 2 Which molecule contains bond angles of both 90° and 120°?
 - \triangle A SF_6
 - \boxtimes **B** PCl₅
 - ☑ C BCl₃
 - **D** BeCl₂

(Total for Question 2 = 1 mark)

- **3** Which molecule has the most polar **bond**?
 - A CH₃CH₂NH₂
 - B CH₃CH₂OH
 - C CH₃CH₂Cl
 - ☑ D CH₃CH₂I

(Total for Question 3 = 1 mark)



 \mathbf{A}

 \square B

 \times C

 \boxtimes D

4 Which is correct for tetrabromomethane, CBr₄?

	Polarity of C-Br bond	Polarity of CBr₄ molecule		
	polar	polar		
⊠ B	non-polar non-polar			
	non-polar polar			
⊠ D	polar	non-polar		

(Total for Question 4 = 1 mark)

5 In alkanes, increasing the length and branching of the carbon chain both affect the boiling temperature.

Which of the following combination of effects is correct?

Effect on boiling temperature		
Increasing chain length	Increasing branching	
increases	decreases	
decreases	increases	
decreases	decreases	
increases	increases	

(Total for Question 5 = 1 mark)

- **6** What is the correct order of boiling temperatures for the hydrogen halides, from the lowest to highest?
 - A HCl, HBr, HI, HF
 - **■ B** HF, HCl, HBr, HI
 - ☑ C HBr, HCl, HF, HI
 - D HI, HBr, HCl, HF

(Total for Question 6 = 1 mark)



- 7 What is **always** formed when s-block nitrates thermally decompose?
 - A A metal nitrite
 - **B** A metal oxide
 - C Nitrogen dioxide
 - D Oxygen

(Total for Question 7 = 1 mark)

- **8** When solid potassium bromide reacts with concentrated sulfuric acid, which substance does **not** form?
 - 🖾 A HBr
 - \square **B** Br₂
 - \boxtimes **C** SO₂
 - ☑ D H₂S

(Total for Question 8 = 1 mark)

- **9** Which equation does **not** represent a disproportionation reaction?
 - \square A Cl_2 + H_2O \rightarrow HClO + HCl
 - \square **B** 3Cl₂ + 6KOH \rightarrow KClO₃ + 5KCl + 3H₂O
 - \square **C** 2KClO₃ \rightarrow 2KCl + 3O₂
 - \square **D** 4KClO₃ \rightarrow 3KClO₄ + KCl

(Total for Question 9 = 1 mark)

- **10** How does the addition of a catalyst to a reaction affect the shape of the Maxwell-Boltzmann distribution curve?
 - A There is no change.
 - **B** The peak moves to the left and is higher.
 - C The peak moves to the right and is lower.
 - D The total area under the curve increases.

(Total for Question 10 = 1 mark)

11 This question is about the equilibrium between sulfur dioxide, oxygen and sulfur trioxide in the gas phase.

$$SO_2(g) + \frac{1}{2}O_2(g) \rightleftharpoons SO_3(g) \Delta H^{\ominus} = -98 \text{ kJ mol}^{-1}$$

(a) What are the effects of **decreasing** the temperature?

(1)

	Effect on rate	Effect on equilibrium yield of SO ₃
⊠ A	increases	decreases
⊠ B	decreases	increases
⊠ C	decreases	decreases
⊠ D	increases	increases

(b) What are the effects of **increasing** the pressure?

(1)

	Effect on rate	Effect on equilibrium yield of SO ₃
⊠ A	increases	decreases
⊠ B	decreases	increases
⊠ C	decreases decreases	
⊠ D	increases	increases

(c) The equation for the reaction can also be written as

$$2SO_2(g) + O_2(g) \rightleftharpoons 2SO_3(g)$$

The enthalpy change for this equation is

(1)

- B -98 kJ mol⁻¹

(Total for Question 11 = 3 marks)



12 A sample of 50 cm³ of ethanol gas is burned completely in 200 cm³ of oxygen.

$$C_2H_5OH(g) + 3O_2(g) \rightarrow 2CO_2(g) + 3H_2O(g)$$

All volumes are measured at a temperature of 400 K and 1 atm pressure.

What is the **total** volume of gas when the reaction is complete?

- \triangle A 150 cm³
- \boxtimes **B** 200 cm³
- \boxtimes **C** 250 cm³
- \square **D** 300 cm³

(Total for Question 12 = 1 mark)

13 Which is a tertiary alcohol?

- A 2-methylbutan-2-ol
- ☑ B 2-methylbutan-1-ol
- C pentan-2-ol
- D pentan-3-ol

(Total for Question 13 = 1 mark)

14 Excess ammonia in ethanol reacts with 1-bromobutane at high pressure.

Which products could be formed in this reaction?

- \square **A** $C_4H_9NH_2$ and NH_4Br
- \square **B** $C_4H_9NH_2$ and C_4H_{10}
- \square **C** C_4H_{10} and HBr
- \square **D** C_4H_8 , NH_4Br and HBr

(Total for Question 14 = 1 mark)

15	-	uestion is about mechanisms and reaction types. e reaction between an alkane and a halogen to form a halogenoalkane is	(1)
	⊠ A	electrophilic addition.	(1)
		nucleophilic addition.	
	⊠ C	free radical substitution.	
	⊠ D	nucleophilic substitution.	
	(b) Th	e reaction between ammonia and a halogenoalkane is	(1)
	⊠ A	electrophilic addition.	
	⊠ B	nucleophilic addition.	
		free radical substitution.	
	⊠ D	nucleophilic substitution.	

(Total for Question 15 = 2 marks)

16	6 Which of these substances causes the greatest amount of anthropogenic global warming?					
	X	A	Carbon dioxide			
	X	В	Methane			
	X	C	Dinitrogen monoxide			
	X	D	Dichlorodifluoromethane			
			(Total for Question 16 = 1 mark)			
17	ln v	whi	ch pair do both substances deplete the ozone layer?			
	X	A	Water vapour and carbon dioxide			
	×	В	Dichlorodifluoromethane and nitrogen monoxide			
	×	C	Dichlorodifluoromethane and carbon dioxide			
	×	D	Water vapour and nitrogen monoxide			
			(Total for Question 17 = 1 mark)			

TOTAL FOR SECTION A = 20 MARKS



SECTION B

Answer ALL the questions. Write your answers in the spaces provided.

- **18** This question is about diamond, graphite and other carbon structures.
 - (a) (i) In diamond, each carbon atom is covalently bonded to four others, in a three-dimensional structure.

Draw a diagram showing this arrangement.

(1)

(ii) Explain the shape and bond angle of this arrangement of carbon atoms in diamond.

(3)

Shape
Bond angle

Explanation



(b)	Graphite consists of a layer lattice, with strong covalent bonds within the layers
	and a weaker force between the layers.

(i)	Draw a diagram to show part of one layer of graphite with between
	13 and 19 carbon atoms, and give the bond angle.

(2)

Diagram

Bond angle	2
------------	---

(ii) Name the force between the layers.

(1)

(iii) Give a reason why graphite conducts electricity whereas diamond does not.

(1)

10



(iv) A graphite shield was used on the front of early spacecraft to preve getting too hot when re-entering the atmosphere.	ent them
Use your knowledge of the structure of graphite and its physical pr suggest two reasons for this use.	roperties to
	(2)
(c) Name one other form of pure carbon.	(1)
(Total for Question	18 = 11 marks)



19	This question is about the structure, properties and reactions of the alcohols, ethano and butan-2-ol.	l
	(a) Ethanol mixes with water in all proportions, but butan-2-ol has limited solubility in water.	
	(i) Name all the intermolecular forces present in these alcohols.	
		(2)
	*(ii) Explain why butan-2-ol has limited solubility in water.	(0)
		(2)
	(b) Both alcohols react with sodium.	
	(i) Describe what you see when ethanol reacts with sodium.	(2)
	(ii) Write the equation for the reaction of ethanol with sodium. State symbols are not required.	(0)
		(2)

(c) E	Both alcohols are separately heated under reflux with acidified	
p	potassium dichromate(VI).	
P	After refluxing, the organic product is distilled from each mixture.	
(i) Name the organic product formed from ethanol and give a chemical test, with the result, to show the functional group present in the product.	(2)
	Name	
Test and	result	
(ii) Give the structure of the organic product formed from butan-2-ol.	(1)
		(1)
(iii) State the ways in which the infrared spectra of butan-2-ol and its oxidation	
	product would differ. Specific wave numbers are not required.	
	specific wave nambers are not required.	(2)
	(Total for Question 19 = 13 ma	rks)
	· · ·	



- **20** This question is about iodine and some of its compounds.
 - (a) Iodine can be obtained from iodine compounds, such as potassium iodide, by reaction with chlorine.
 - (i) Write the **ionic** equation for the formation of iodine, by the addition of chlorine to aqueous potassium iodide.

 State symbols are not required.

(1)

(ii) The colour of the iodine solution formed is red-brown.

Name an organic solvent that can be added to extract iodine from its aqueous solution, and give the colour of the organic layer.

(2)

Name

Colour

(iii) lodine reacts with sodium thiosulfate solution.

$$I_2(aq) + 2Na_2S_2O_3(aq) \rightarrow 2NaI(aq) + Na_2S_4O_6(aq)$$

State which element is oxidised and which is reduced, giving the relevant changes in oxidation number.

(2)



(i) Draw	the skeleta	I formulae of	these two i	somers.		(1)
using Sugg	mass spect est why 1-io	easily distingu crometry beca odopropane of and 1-bromo	use they on nly has one	lly give a sing molecular ion	le molecular i n peak wherea	on peak. as
		£ 1 :		a noak at m/	e = 43.	
	nass spectru	e of the ion re	·			it is formed. (2)
Give 1-iodopr hot silve	opane and 2	•	sponsible fo	tely dissolved	nd state how	(2)
Give 1-iodopr hot silve (i) State	opane and 2 r nitrate solu	e of the ion re	e are separa A precipita	tely dissolved ate forms in e	I in ethanol ar	(2) and



(d) Aqueous sodium hydroxide reacts with 1-iodopropane to form propan-1-ol.

Draw the mechanism for this reaction. Include curly arrows and relevant lone pairs.

(2)

- (e) Hot concentrated potassium hydroxide in ethanol reacts with 1-iodopropane and 2-iodopropane.The same organic compound is formed in each case.
 - (i) State the type of reaction occurring.

(1)

(ii) **Name** the organic product formed in both reactions.

(1)

(Total for Question 20 = 16 marks)

TOTAL FOR SECTION B = 40 MARKS

SECTION C

Answer ALL the questions. Write your answers in the spaces provided.

21 'Hard' water is water that contains significant concentrations of calcium ions or magnesium ions.

Hard water may be produced when rainwater, containing dissolved carbon dioxide, passes through rocks containing calcium carbonate. Calcium hydrogencarbonate is formed in solution.

Hard water may also be formed when rain water passes through minerals containing calcium sulfate or magnesium sulfate.

Hard water causes two problems.

The first is that it forms a precipitate (scum) when mixed with soap.

The second is that, on heating, it will form a deposit of calcium or magnesium carbonate which reduces the efficiency of heating elements and may lead to blockages in boiler pipes.

- (a) A flame test can be used to detect the metal ion present in the precipitate formed by heating hard water.
- *(i) Explain how metal ions produce a colour in a flame test.

 (3)

 (ii) Give the colour of the flame you would expect to see when calcium ions are present.

 (1)

 (iii) Give the reason why magnesium ions do not produce a flame colour.

 (1)



(b) Write an equation to show how calcium hydrogencarbonate is formed when rainwater, containing dissolved carbon dioxide, reacts with calcium carbonate. State symbols are not required.

(1)

(c) Suggest why calcium sulfate and magnesium sulfate can form hard water but barium sulfate cannot.

(1)

(d) Group 2 metal carbonates decompose on heating.

$$MCO_3 \rightarrow MO + CO_2$$

*(i) Explain why calcium carbonate requires stronger heating to decompose than magnesium carbonate.

(3)

 •••••			

(ii) When 10.00 g of a Group 2 metal carbonate is completely decomposed by heating, 1.626 dm³ of carbon dioxide forms, at room temperature and pressure (r.t.p.).

Deduce by calculation the metal ion present.

[Molar volume of a gas at r.t.p. = $24.0 \,\mathrm{dm^3 \,mol^{-1}}$]

(3)

(iii) Calcium oxide dissolves in water to form calcium hydroxide (limewater), ${\rm Ca(OH)_2(aq)}$.

Write the equation for the reaction of carbon dioxide with limewater. Include state symbols.

(1)

(e) The solubility of calcium hydroxide in water is determined by titration of a saturated solution.

An excess of calcium hydroxide is added to about 100 cm³ of distilled water in a conical flask.

The flask is stoppered, shaken and allowed to stand overnight.

10.0 cm³ portions of this saturated solution are titrated with 0.0500 mol dm⁻³ hydrochloric acid.

$$\mathrm{Ca(OH)_2} \ + \ \mathrm{2HCl} \ \rightarrow \ \mathrm{CaCl_2} \ + \ \mathrm{H_2O}$$

(i) Suggest a suitable indicator for this titration and give the colour change you would expect to see at the end-point.

(2)

Indicator

Colour change from ______ to ____

(ii) The mean titre is 8.90 cm³.

Calculate the concentration of the saturated calcium hydroxide solution, in g dm⁻³.

(4)

(Total for Question 21 = 20 marks)

TOTAL FOR SECTION C = 20 MARKS TOTAL FOR PAPER = 80 MARKS



BLANK PAGE

BLANK PAGE



The Periodic Table of Elements

0 (8)	(18)	4.0	H	helium 2	20.2	Š	neon	10
7				(17)	19.0	L	fluorine	6
9				(16)	16.0	0	oxygen	œ
2				(15)	14.0	z	nitrogen	7
4				(14)	12.0	U	carbon	9
ъ				(13)	10.8	В	poron	2
		0:1	L phydosen	1				
					_			
				Key	relative atomic mass	atomic symbol	name	atomic (proton) number
2				i) Key	.0 relative atomic mass	то	llium	4 atomic (proton) number
2				(2) Key	9.0 relative atomic mass	Be atomic symbol	n beryllium	4 atomic (proton) number

	Elements with atomic numbers 112-116 have been reported	but not fully authenticated
[272]	Sg	roentgenium
[271]	S	lamstadtium

Ξ

darmstadtiu 110 õ

neitnerium

[277] **Hs** hassium 108

[264] **Bh** bohrium 107

Sg seaborgium 106

[262] **Db**dubnium

בּ

Ac* actinium

Radium

Fr francium

[261]

[227]

[226]

[223]

105

5

8

88

109

[268]

83.8 Krypton 36

Br bromine

selenium

arsenic

ermaniun

32

31

Se 34

As 33

g

Ga gallium

Znc zinc 30

63.5 copper 29

58.7 **Bi** nickel 28

58.9 Cobalt 27

55.8 Fe iron 26

Mn manganese

chromium

vanadium

T

scandium

calcium

S

K

23

22

21

20

19

24

54.9

52.0

47.9

45.0

40.1

39.1

12

Ξ

79.9

79.0

74.9 15

72.6

69.7 13

17

Ar argon 18

35.5 CL chlorine

∨lfu

surodphorus

Silicon 14

Al

(12)

6

0

9

(2)

£

Mg magnesium

23.0 Na sodium

24.3

28.1

131.3 Xenon xenon

126.9

127.6

118.7

114.8

112.4

107.9

106.4

[222] **Rn** radon 86

astatine

8

¥

[209] **Po** polonium

209.0 **Bi** bismuth 83

207.2 **Pb** tead 82

204.4 Tl thallium

200.6

197.0

192.2

190.2

186.2

183.8

180.9

178.5

137.3

5

4

43

42

4

6

yttrium 39 138.9

38

37

8

Hg mercury 80

Au gold 79

Pt Pt platinum 78

Ir iridium 77

Os osmium 76

Re

≥

Тa

Ŧ

La*

Ba barium

CS Caesium

tungsten 74

tantalum 73

hafnium 72

anthanum

57

26

55

24

[210]

iodine

tellurium

52

ъ

121.8 Sb antimony 51

I indium

Cadmium

Ag Silver

Pd palladium 46

Rh rhodium

Tc Ru ruthenium ruthenium

95.9 Mo

Nobium miobium

Zr zirconium

Sr strontium

rubidium

92.9

91.2

88.9

87.6

85.5 **Rb**

4

48

	140	141		[147]	150	152	157	159	163	165	167	169	173	175
series	ő	ڇ	Ž	Pa	Sm	En	В	4	ð	운	Б	Ē	χ	3
ioe	cerium	praecodymium	ĕ	promethium	samarium	europium	gadolinium	terbium	dysprosium	holmium	erbium	thulium	ytterbium	lutetium
5	28	29		19	62	63	64	65	99	29	89	69	70	71
	232	232 [231]	11	[237]	[242]	[243]	[247]	[245]	[251]	[254]	[253]	[326]	[254]	[257]
	두	Pa	_	2	Pn	Am	క	쓢	້ຽ	ß	F	PW	ž	۲
	thorium	protactinium	3	neptunium	plutonium	americium	arrium	berkelium	californium	einsteinium	fermium	mendelevium	nobelium	lawrencium
	8	91		93	4	92	%	46	86	66	100	101	102	103

Lanthanide se

Actinide serie