Please check the examination details below before entering your candidate information				
Candidate surname	Other names			
Pearson Edexcel International Advanced Level	e Number Candidate Number			
Friday 18 January 2019				
Afternoon (Time: 1 hour 15 minutes)	Paper Reference WCH03/01			
Chemistry Advanced				
Unit 3: Chemistry Laboratory Skills I				
Candidates must have: Scientific calc	Total Marks			

# Instructions

- Use **black** ink or **black** ball-point pen.
- Fill in the boxes at the top of this page with your name, centre number and candidate number.
- Answer **all** questions.
- Answer the questions in the spaces provided
  - there may be more space than you need.

# Information

- The total mark for this paper is 50.
- The marks for **each** question are shown in brackets
  - use this as a guide as to how much time to spend on each question.
- You will be assessed on your ability to organise and present information, ideas, descriptions and arguments clearly and logically, including your use of grammar, punctuation and spelling.
- A Periodic Table is printed on the back cover of this paper.

## **Advice**

- Read each question carefully before you start to answer it.
- Show all your working in calculations and include units where appropriate.
- Check your answers if you have time at the end.

Turn over ▶





# Answer ALL the questions. Write your answers in the spaces provided.

- 1 A white solid A contains one cation and one anion.
  - (a) A small amount of solid **A** was placed in a test tube and aqueous sodium hydroxide added. The mixture was warmed gently.

Complete the inference column in the table.

(2)

Observation	Inference
A pungent smelling gas was evolved that turned damp red litmus paper blue	The gas formed is
	The <b>formula</b> of the cation in <b>A</b> is

(b) (i) An aqueous solution of **A** was placed in a test tube and acidified with dilute nitric acid. A few drops of silver nitrate solution were added. Complete the inference column in the table.

(1)

Observation	Inference
Cream precipitate formed	The precipitate is

(ii) Write the **ionic** equation, including state symbols, for the formation of the cream precipitate in (b)(i).

(2)



(iii) Describe how you would confirm the identity of the <b>anion</b> in the cream precipitate formed in (b)(i).		
ioimed iii (b)(i).	(2)	
	(Total for Question 1 = 7 marks)	

**2** (a) A student was provided with aqueous solutions of four compounds:

barium nitrate

hydrochloric acid

sodium carbonate

sulfuric acid

Four bottles, labelled **B**, **C**, **D** and **E**, each contained one of the solutions. The student mixed pairs of the solutions to determine which solution was in each bottle.

The results are shown.

Solutions mixed	Observations		
<b>B</b> and <b>C</b>	Effervescence with bubbles of a colourless gas given off		
<b>B</b> and <b>D</b>	No visible change		
<b>B</b> and <b>E</b>	A white precipitate formed which did <b>not</b> dissolve on the addition of dilute nitric acid		
<b>C</b> and <b>D</b>	Effervescence with bubbles of a colourless gas given off		
<b>C</b> and <b>E</b>	A white precipitate formed which dissolved with effervescence on the addition of dilute nitric acid		
<b>D</b> and <b>E</b>	No visible change		

Use the observations in the table to deduce the identity of the compound in each bottle. Identify each compound by name or formula.

В	
C	
D	
_	

(3)



(b) (i)	The identity of the <b>cations</b> present in barium nitrate and sodium carbonate can be confirmed with a flame test on the solid compounds.	
	Describe how you would carry out a flame test.	(3)
(ii)	State the flame colours produced by barium nitrate and sodium carbonate.	
	Barium nitrate	
	Sodium carbonate	(2)
	(Total for Question 2 = 8 m	arks)



3 Chlorine-based bleaches contain sodium chlorate(I), NaClO, as the active ingredient. The concentration of NaClO in bleach was determined by a titration method using sodium thiosulfate.

Sodium chlorate(I) reacted with potassium iodide in acidic solution to produce iodine.

$$ClO^{-} + 2I^{-} + 2H^{+} \rightarrow I_{2} + Cl^{-} + H_{2}O$$

The iodine was then titrated with sodium thiosulfate.

$$2S_2O_3^{2-} + I_2 \rightarrow 2I^- + S_4O_6^{2-}$$

### **Procedure**

- 1. A burette was filled with 0.0600 mol dm<sup>-3</sup> sodium thiosulfate solution.
- 2. 10.0 cm³ of bleach was pipetted into a 250.0 cm³ volumetric flask and excess potassium iodide and sulfuric acid were added to release iodine. The volume was made up to the mark with distilled water.
- 3. 25.0 cm<sup>3</sup> of this solution was pipetted into a conical flask and titrated with the sodium thiosulfate solution using a suitable indicator.
- (a) State the indicator used and give the colour change at the end-point.

(2)

Indicator	Colour change at the end-point	
	From	to

(b) (i) Complete the table of results.

(1)

Number of titration	1	2	3	4
Burette reading (final) / cm <sup>3</sup>	23.65	46.45	24.40	47.10
Burette reading (start) / cm <sup>3</sup>	0.00	23.65	1.20	24.40
Titre / cm³				

(ii) State with a reason which results should be used to calculate the mean titre value.

(2)

(iii) Calculate the mean titre.

(1)

(iv) Calculate the number of moles of sodium thiosulfate in this mean titre.

(1)

(v) Calculate the number of moles of iodine in 25.0 cm<sup>3</sup> of the diluted solution.

(1)

(vi) Calculate the number of moles of sodium chlorate(I) in the 250.0 cm<sup>3</sup> volumetric flask.

(1)

(vii) Calculate the concentration of sodium chlorate(I) in the **undiluted** bleach in mol dm<sup>-3</sup>.

(1)



	(Total for Question 3 = 13 ma	rks)
	No calculations are required.	(3)
	Describe the steps you would take to prepare this standard solution as accurately as possible. You are supplied with the appropriate mass of sodium thiosulfate and the usual laboratory glassware, including a volumetric flask.	
(c)	The 0.0600 mol dm <sup>-3</sup> sodium thiosulfate solution used in this titration is known as a standard solution.	

**BLANK PAGE** 




4 Hydrogen peroxide, H<sub>2</sub>O<sub>2</sub>, decomposes according to the equation

$$2H_2O_2 \rightarrow 2H_2O + O_2$$

The rate of decomposition is increased by a catalyst.

A student tested three metal oxides to determine which was the best catalyst. The oxides were manganese(IV) oxide, iron(III) oxide and lead(IV) oxide. They are all solids.

The student used the following apparatus and experimental procedure.



# **Procedure**


- 1. Hydrogen peroxide solution was poured into the conical flask.
- 2. Solid manganese(IV) oxide was added.
- 3. The bung was quickly replaced to connect the gas syringe to the conical flask.
- 4. The procedure was repeated using iron(III) oxide and lead(IV) oxide.

when using this procedure.	(3)
) State the measurements the student should make to determine which is the best catalyst.	
Sest eatalyst.	(2)
The student thought that some of the gas escaped from the conical flask before	
the bung had been replaced.	
Suggest how this experiment could be modified to prevent this loss.	
	(1)



(d) Another student thought that some of the oxygen produced may have come from the decomposition of the metal oxide.	
Suggest how this idea could be tested.	(2)
(Total for Question 4 = 8 mark	ks)

**5** Some organic reactions are shown.



(a) Reaction 1 and Reaction 2 use the same reagent but require different conditions. Identify the reagent and give the conditions needed for Reaction 1.

(2)

(b) (i) Give a chemical test and its positive result to show the presence of the double bond in  $CH_3CH = CH_2$ .

(2)

(ii) Give the structure of the organic product of the test in (b)(i).

(1)



CH <sub>3</sub> CH <sub>2</sub> CH <sub>2</sub> OI	ded phosphorus(V) chloride, PCl₅, to the H. Hydrogen chloride was formed. observation the student would be expec	•	(1)
for each o	the table to show the hazard and the ap chemical. clude the wearing of eye protection and		(3)
Chemical	Hazard	Safety precaution	n
PCl₅			
CH <sub>3</sub> CH <sub>2</sub> CH <sub>2</sub> OH			
HCl			
potassium di	, CH₃CH₂CH₂OH is oxidised to CH₃CH₂CH0 chromate(VI) acidified with sulfuric acid. colour <b>change</b> that occurs during this ox		
(i) State the	colour <b>change</b> that occurs during this ox	duation reaction.	(1)

(ii) Draw a labelled diagram of the apparatus you would use to carry out Reaction **3** and collect the product.

(3)

(iii) Explain how infrared spectroscopy could be used to confirm that **all** the CH<sub>3</sub>CH<sub>2</sub>CH<sub>2</sub>OH has been oxidised to CH<sub>3</sub>CH<sub>2</sub>CHO in Reaction **3**. You are not expected to give specific wavenumbers.

(1)

(Total for Question 5 = 14 marks)

**TOTAL FOR PAPER = 50 MARKS** 



# The Periodic Table of Elements

1		
9		
2		
_		
272		
4		
m		
	1.0	
	1.0	
	1.0	
2		
2		
2		
1 2		

_							1
0 (8) (18) 4.0	He hetium 2	20.2 Ne neon	39.9 Ar argon 18	83.8 Krypton 36	131.3 Xe xenon 54	[222] Rn radon 86	ted
7	(17)	19.0 F fluorine 9	35.5 Cl chlorine 17	79.9 Br bromine 35	126.9 I lodine 53	[210] At astatine 85	een repo
9	(16)	16.0 O oxygen 8	32.1 <b>S</b> sulfur 16	Se selenium 34	127.6 Te tellurium 52	Po polonium 84	116 have t
10	(15)	14.0 N nitrogen 7	31.0 Phosphorus 15	As As arsenic 33	Sb antimony 51	209.0 Bi bismuth 83	tomic numbers 112-116 hav but not fully authenticated
4	(14)	12.0 C carbon 6	Si silicon	72.6 Ge germanium 32	118.7 Sn tin 50	207.2 <b>Pb</b> tead 82	atomic nun but not fu
m	(13)	10.8 <b>B</b> boron 5	27.0 AI aluminium 13	Ga gailtium 31	In In indium 49	204.4 Tl thallium 81	Elements with atomic numbers 112-116 have been reported but not fully authenticated
			(12)	<b>Zn</b> zinc 30	Cd Cd cadmium 48	Hg mercury 80	Elem
			(11)	63.5 Cu copper 29	Ag silver 47	197.0 <b>Au</b> gold 79	Rg coentgenium 111
			(01)	58.7 Ni nickel 28	Pd Pd palladfum 46	Pt platinum 78	Ds damstadtum r 110
			(6)	S8.9 Co cobalt 27	Rh rhodium 45	192.2 Ir iridium 77	[268] Mt meltnerium 109
0. =	hydrogen 1		(8)	55.8 Fe iron 26	Ru ruthenium 44	Os Osmium 76	Hs Hassium r 108
			0	Mn Manganese 25	_ §	Re rhenium 75	[264] <b>Bh</b> bohrium 107
		mass ool umber	(9)	52.0 54.9 Cr Mn chromium manganese 24 25	Mo molybdenum 42	183.8 W tungsten 74	Sg seaborgium 106
	Key	relative atomic mass atomic symbol name atomic (proton) number	(5)	50.9 V vanadium 23	92.9 Nb niobium 41	180.9 Ta tantalum 73	[262] <b>Db</b> dubnium 105
		relativ atoric atomic	(4)	47.9 Ti titanium 22	91.2 Zr zirconium 40	Hf Hafnium 72	[261] Rf nutherfordium 104
			(3)	Sc scandium 21	88.9 <b>Y</b> yttrium 39	La* La* Lanthanum 57	Ac* actinium 89
7	(2)	9.0 Be beryttium	Mg magnesium 12	Ca calcium 20	87.6 Sr strontium 38	137.3 Ba barium 1 56	[226] Ra radium 88
-	ε	6.9 Li lithium 3	Na sodium	39.1 <b>K</b> potassium 19	Rb rubidium 37	CS Caesium 55	[223] Fr francium 87

Lanthanide series	
anthanid	S
anthanid	er
anthani	9
anthar	ğ
an	ᇹ
B	Ę
	B

<sup>\*</sup> Actinide series

_	141	144	[147]	150	152	157	159	163	165	167	169	173	175
	P.	P	Pm	Sm	Eu	В	T <sub>P</sub>	δ	유	ᆸ	Ē	χ	ב
E	praseodymium	neodymium	promethium	samarium	europium	gadolinium	terbium	dysprosium	holmium	erbium	thulium	ytterbium	lutetium
28	59	9	61	62	63	64	9	99	29	89	69	70	71
2	[231]	238	[237]	[242]	[243]	[247]	[245]	[251]	[254]	[253]	[256]	[254]	[257]
£	Pa	_	å	Pa	Am	5	BK	ຽ	Es	Fm	PW	9 N	۲
Eni	protactinium	uranium	neptunium	plutonium	americium	curium	berkelium	californium	einsteinium	fermium	mendelevium	nobelium	lawrencium
0	16	92	93	94	95	96	46	86	66	100	101	102	103