



Pearson
Edexcel

Mark Scheme (Results)

January 2022

Pearson Edexcel International A Level
in Statistics S2 (WST02) Paper 01

Edexcel and BTEC Qualifications

Edexcel and BTEC qualifications are awarded by Pearson, the UK's largest awarding body. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information visit our qualifications websites at www.edexcel.com or www.btec.co.uk. Alternatively, you can get in touch with us using the details on our contact us page at www.edexcel.com/contactus.

Pearson: helping people progress, everywhere

Pearson aspires to be the world's leading learning company. Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: www.pearson.com/uk

January 2022

Question Paper Log Number P71201A

Publications Code WST02_01_2201_MS

All the material in this publication is copyright

© Pearson Education Ltd 2022

General Marking Guidance

- All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.
- Mark schemes should be applied positively. Candidates must be rewarded for what they have shown they can do rather than penalised for omissions.
- Examiners should mark according to the mark scheme not according to their perception of where the grade boundaries may lie.
- There is no ceiling on achievement. All marks on the mark scheme should be used appropriately.
- All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e. if the answer matches the mark scheme. Examiners should also be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.
- Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.
- When examiners are in doubt regarding the application of the mark scheme to a candidate's response, the team leader must be consulted.
- Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.

EDEXCEL IAL MATHEMATICS

General Marking Guidance

- All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.
- Mark schemes should be applied positively. Candidates must be rewarded for what they have shown they can do rather than penalised for omissions.
- Examiners should mark according to the mark scheme not according to their perception of where the grade boundaries may lie.
- There is no ceiling on achievement. All marks on the mark scheme should be used appropriately.
- All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e. if the answer matches the mark scheme. Examiners should also be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.
- Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.
- When examiners are in doubt regarding the application of the mark scheme to a candidate's response, the team leader must be consulted.
- Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.

General Instructions for Marking

1. The total number of marks for the paper is 75.
2. The Edexcel Mathematics mark schemes use the following types of marks:
 - **M** marks: method marks are awarded for 'knowing a method and attempting to apply it', unless otherwise indicated.
 - **A** marks: Accuracy marks can only be awarded if the relevant method (M) marks have been earned.
 - **B** marks are unconditional accuracy marks (independent of M marks)
 - Marks should not be subdivided.
3. Abbreviations


These are some of the traditional marking abbreviations that will appear in the mark schemes.

- bod – benefit of doubt
 - ft – follow through
 - the symbol \square will be used for correct ft
 - cao – correct answer only
 - cso - correct solution only. There must be no errors in this part of the question to obtain this mark
 - isw – ignore subsequent working
 - awrt – answers which round to
 - SC: special case
 - oe – or equivalent (and appropriate)
 - dep – dependent
 - indep – independent
 - dp decimal places
 - sf significant figures
 - * The answer is printed on the paper
 - \square The second mark is dependent on gaining the first mark
4. All A marks are 'correct answer only' (cao.), unless shown, for example, as A1 ft to indicate that previous wrong working is to be followed through. After a misread however, the subsequent A marks affected are treated as A ft, but manifestly absurd answers should never be awarded A marks.
 5. For misreading which does not alter the character of a question or materially simplify it, deduct two from any A or B marks gained, in that part of the question affected.
 6. If a candidate makes more than one attempt at any question:
 - If all but one attempt is crossed out, mark the attempt which is NOT crossed out.
 - If either all attempts are crossed out or none are crossed out, mark all the attempts and score the highest single attempt.
 7. Ignore wrong working or incorrect statements following a correct answer
Special notes for marking Statistics exams (for AAs only)
 - Any correct method should gain credit. If you cannot see how to apply the mark scheme but believe the method to be correct then please send to review.
 - For method marks, we generally allow or condone a slip or transcription error if these are seen in an expression. We do not, however, condone or allow these errors in accuracy marks.

Question Number	Scheme		Marks
1 (a)	$X = \text{faults in a week} \Rightarrow X \sim \text{Po}(6)$		
	$[P(X \geq x) = 0.1528 \Rightarrow P(X \leq x - 1)] = 0.8472$		M1
	Using tables $P(X \leq 8) = 0.8472 \Rightarrow x - 1 = 8$		M1
	$x = 9$		A1
			(3)
(b)	$Y = \text{faults in six weeks} \Rightarrow Y \sim N(36, 36)$		B1
	$P(Y < 32) = P\left(Z < \frac{31.5 - 36}{6}\right) [= P(Z < -0.75)]$		M1 M1
	$= 0.2266$	awrt 0.227	A1
(c)	$W = \text{Number of poor weeks} \Rightarrow W \sim B(50, 0.1528)$		B1
	$[P(W > 1)] = 1 - P(W \leq 1)$		M1
	$= 1 - (0.8472^{50} + 50 \times 0.1528 \times 0.8472^{49})$		dM1
	$= 0.99748\dots$	awrt 0.997	A1
			(4)
Notes			Total 11
1 (a)	M1	Writing or using $1 - P(X \leq x - 1)$	
	M1	For 0.8472 May be implied by $x - 1 = 8$	
	A1	$x = 9$	
(b)	B1	Writing or using $N(36, 36)$ (May be implied by a correct standardisation expression)	
	M1	Standardising with 30.5/31/31.5/32/32.5/39.5/40/40.5/41/41.5, their mean and standard deviation (Allow \pm)	
	M1	A fully correct standardisation. May be implied by ± 0.75	
	A1	awrt 0.227	
(c)	B1	Writing or using $B(50, 0.1528)$	
	M1	Writing or using $1 - P(W \leq 1)$ (Allow any letter)	
	dM1	Dependent on using binomial. Using $1 - [P(W = 0) + P(W = 1)]$ (implied by awrt 0.997 or 0.9975 or $1 - \text{awrt } 0.00257$) Using binomial may be implied by $(1 - p)^{50} + {}^n C_r \times p \times (1 - p)^{49}$ where p is a probability Condone ${}^n C_r$ missing	
	A1	awrt 0.997 or 0.9975	

Question Number	Scheme		Marks	
2 (a)	$f(x) = \begin{cases} \frac{1}{4k} & -k \leq x \leq 3k \\ 0 & \text{otherwise} \end{cases}$		M1 A1	
			(2)	
(b)	$[E(X)] = k$		B1	
			(1)	
(c)	$[\text{Var}(X)] = \frac{(3k - -k)^2}{12} = \frac{16k^2}{12} \quad \text{or} \quad \left[\frac{x^3}{3} "f(x)" \right]_{-k}^{3k} - ("k")^2$		M1	
			$= \frac{4k^2}{3} *$	A1* cso
			(2)	
(d)	$E(X^2) = \text{Var}(X) + E(X)^2 = \frac{4k^2}{3} + ("k")^2$		M1	
			$= \frac{7k^2}{3}$	A1
			$E(3X^2) = 3E(X^2) = 3 \times \frac{7k^2}{3} = 7k^2$	A1
			(3)	
Notes			Total 8	
2 (a)	M1	For the 1 st line of the pdf including the inequality, allow use of < instead of one/both ≤ signs		
	A1	Fully correct, allow use of < instead of one/both ≤ signs. Allow equivalent for the 0 otherwise.		
(b)	B1	Cao		
(c)	M1	Use of $\text{Var}(X) = \frac{(\beta - \alpha)^2}{12}$ or $\left[\frac{x^3}{3} "f(x)" \right]_{-k}^{3k} - ("k")^2$		
			A1* cso	Answer is given. Correct solution only with no incorrect working.
(d)	M1	Use of $E(X^2) = \text{Var}(X) + E(X)^2$ ft their $E(X)$ or $\left[\frac{x^3}{3} "f(x)" \right]_{-k}^{3k}$ this integration may be seen in part (c) or part (d)		
			A1	$\frac{7k^2}{3}$ (This must be seen in part (d)) May be implied by $7k^2$)
	A1	Cao		

Question Number	Scheme		Marks
3 (a)	We can assume breakdowns are [rare], independent events occurring at a constant rate.		B1 (1)
(b)	$H_0 : \lambda = 8$ $H_1 : \lambda \neq 8$		B1 (1)
(c)	$X \sim \text{Po}(8)$		
	$P(X \leq 2) = 0.0138$ oe $P(X \leq 3) = 0.0424$ oe		M1
	$P(X \geq 14) = 0.0342$ oe $P(X \geq 15) = 0.0173$ oe		M1
	$X \leq 2 \cup X \geq 15$ oe		A1 (3)
(d)	“0.0138” + “0.0173”		M1
	=“0.0311”		A1ft (2)
(e)	“[4 is] not in the critical region”		M1
	So there is insufficient evidence that refurbishment has changed the mean breakdown rate		A1 (2)
Notes			Total 9
3 (a)	B1	A correct statement which include the words independent or constant rate or singly. No context needed	
(b)	B1	Both hypotheses correct. Must be attached to H_0 and H_1 in terms of λ or μ .	
(c)	M1	Use of Po(8) to find the lower critical value. May be implied by either 0.0138 or 0.0424 or $X \leq 2$ if no probabilities shown (Calculator values: 0.01375... and 0.04238...)	
	M1	Use of Po(8) to find the upper critical value. May be implied by 0.0342 or 0.0173 or 0.9658 or 0.9827 or $X \geq 15$ if no probabilities shown (Calculator values: 0.03418... and 0.01725... and 0.96581... and 0.98274...)	
	A1	$X \leq 2$ oe $[\cup] X \geq 15$ oe Condone the use of and/or Do not allow as probability statements Allow $[0, 2]$ or $[0, 3)$ and $[15, \infty]$ or $[15, \infty)$ or $(14, \infty]$ or $(14, \infty)$	
(d)	M1	Adding the two probabilities for their critical region	
	A1ft	0.0311 Allow 3.11 or awrt 3.1[0] or awrt 0.031[0] ft their critical region	
		NB 3.11 or 0.0311 or awrt 3.1[0] or awrt 0.031[0] will score 2/2	
(e)	M1	A correct statement ft their critical region e.g. Do not reject H_0 /Accept H_0 /not significant – no context needed but do not allow contradicting non contextual comments	
	A1	Correct conclusion in context. Must include rate/number of breakdown (Allow decreased for changed)	
		NB Award M1 A1 for a correct contextual statement on its own	

Question Number	Scheme		Marks
4 (a)			B1 B1
			(2)
	(b)	$\frac{1}{2}(3+9) \times k = 1$ or $\frac{1}{2}(3-1)k + (6-3)k + \frac{1}{2}(10-6)k = 1$ or $\frac{1}{2}k \left[\frac{x^2}{2} - x \right]_1^3 + k[x]_3^6 + \frac{1}{4}k \left[10x - \frac{x^2}{2} \right]_6^{10} = 1$	M1
		$k = \frac{1}{6} *$	A1* cso
		(2)	
(c)	$\int_1^x \frac{1}{12}(x-1) dx$ or $\int \frac{1}{12}(x-1) dx$ and using $F(1) = 0$		M1
	$\int_3^x \frac{1}{6} dx + "F(3)"$ or $\int \frac{1}{6} dx$ and using " $F(3) = \frac{1}{6}$ "		M1
	$\int_6^x \left(\frac{5}{12} - \frac{1}{24}x \right) dx + "F(6)"$ or $\int \left(\frac{5}{12} - \frac{1}{24}x \right) dx$ and using either " $F(6) = \frac{2}{3}$ " or $F(10) = 1$		M1
	$F(x) = \begin{cases} 0 & x < 1 \\ \frac{1}{24}(x^2 - 2x + 1) & 1 \leq x \leq 3 \\ \frac{1}{6}(x-2) & 3 < x \leq 6 \\ \frac{1}{48}(20x - x^2 - 52) \text{ or } 1 - \frac{(10-x)^2}{48} & 6 < x \leq 10 \\ 1 & x > 10 \end{cases}$		A1oe A1oe A1 oe B1
		(7)	
(d)	$P(X > E(X)) = 1 - F\left(\frac{61}{12}\right) = 1 - 0.51388... = 0.4861...$	awrt 0.486	M1 A1 (2)
(e)	Since (d) < 0.5 [the mean is greater than the median] therefore positive (skew) or follow through their sketch in part (a)		M1 A1ft (2)
Notes			Total 15
4(a)	B1	Correct shape. Must start and end on the x axis	
	B1	Fully correct including 1, 3, 6, 10 and k . Allow $\frac{1}{6}$ for k Ignore labels for x and $f(x)$ and any extras e.g. $k/2$	
(b)	M1	Setting up the area of the trapezium = 1 or 2 triangles + a rectangle = 1 or a fully correct integration, including limits = 1	
	A1* cso	Answer is given. Correct solution only with no incorrect working.	
(c)	M1	For a correct method to find the 2 nd line Allow in terms of k	

	M1	For a correct method to find the 3 rd line, ft their $F(3)$. If using + c method then ft their $F(3) = \frac{1}{6}$ Allow in terms of k
	M1	For a correct method to find the 4 th line, ft their $F(6)$. If using + c method then ft their $F(6) = \frac{2}{3}$ Allow in terms of k
	A1	2 nd line correct including inequality. Allow $<$ instead of \leq
	A1	3 rd line correct including inequality. Allow $<$ instead of \leq
	A1	4 th line correct including inequality. Allow $<$ instead of \leq
	B1	1 st and 5 th line correct. Allow “otherwise” for the range on the 1 st or 5 th line but not both. All 5 lines must be in terms of the same letter.
(d)	M1	For use of $1 - F\left(\frac{61}{12}\right)$ using the their line of $F(x)$ for $3 < x \leq 6$. May use integration/area methods
	A1	awrt 0.486 Allow $\frac{35}{72}$
(e)	M1	For correctly comparing part (d) with 0.5 (may be implied by a correct comparison of mean and median (5)) do not allow mean is greater than the median on its own
	A1ft	For positive skew or ft their answer to part (d) Accept “no (or negligible) skew” following a reason that “mean \approx median” Allow argument based on sketch in part (a)

Question Number	Scheme		Marks
5 (a)	B(n , 0.045)		B1 (1)
(b)	<u>Applicants</u> are independent (no identical twins) or the <u>proportion/probability</u> identified as <u>colour blind</u> does not change over time		B1 (1)
(c)	B(120, 0.045) \Rightarrow Po(5.4)		B1
	$P(X = 5) = \frac{e^{-5.4} \times 5.4^5}{5!}$		M1
	= 0.1728... awrt 0.173		A1 (3)
(d)	Binomial with large n and very small p		B1 B1 (2)
	$H_0 : p = 0.75 \quad H_1 : p \neq 0.75$ B(96, 0.75) \Rightarrow N(72, 18)		B1 B1
	$Z = \frac{67.5 - 72}{\sqrt{18}} \quad \text{or} \quad \frac{x \pm 0.5 - 72}{\sqrt{18}}$ = -1.06066... or $\frac{x + 0.5 - 72}{\sqrt{18}} < -1.96$ or $\frac{x - 0.5 - 72}{\sqrt{18}} > 1.96$ P($z < -1.06$) = 0.1444... / 0.1446 or CR < 63.2 awrt 0.144 or 0.145 There is insufficient evidence to reject H_0 Insufficient evidence against Jaymini's claim		M1 A1 A1 dM1 A1 (7)
ALT	Let p be the probability of an applicant fail to become a pilot. $H_0 : p = 0.25 \quad H_1 : p \neq 0.25$ B(96, 0.25) \Rightarrow N(24, 18)		B1 B1
	$Z = \frac{28.5 - 24}{\sqrt{18}} \quad \text{or} \quad \frac{x \pm 0.5 - 24}{\sqrt{18}}$ = 1.06066... or $\frac{x + 0.5 - 24}{\sqrt{18}} < -1.96$ or $\frac{x - 0.5 - 24}{\sqrt{18}} > 1.96$ P($z > 1.06$) = 0.1444... / 0.1446 or CR > 32.8 awrt 0.144 or 0.145 There is insufficient evidence to reject H_0 Insufficient evidence against Jaymini's claim		M1 A1 A1 dM1 A1 (7)
Notes			Total 14
5 (a)	B1	For binomial with correct parameters n and 0.045	
(b)	B1	For one of the given reasons. Must have context Allow equivalent statements Do not allow number for proportion/probability	
(c)	B1	Using or writing Po(5.4)	
	M1	For $\frac{e^{-\lambda} \lambda^5}{5!}$ with any value for λ	
	A1	awrt 0.173	
		NB A correct answer with no incorrect working scores 3/3	
(d)	B1	n is large (Allow number of trials for n)	

	B1	p is small (Allow probability for p)
(e)	B1	Both hypotheses correct in terms of p or π Must be attached to H_0 and H_1
	B1	For writing or using $N(72, 18)$ (May be implied by a correct standardisation expression)
	M1	Standardising using 67.5 or 67 or 66.5 or $x \pm 0.5$ with their mean and standard deviation (Allow \pm)
	A1	awrt -1.06 (may be implied by awrt 0.144 or 0.145) or a correct standardisation with ± 1.96 (ignore incorrect inequality symbol and allow =)
	A1	Using a probability route: awrt 0.144 or 0.145 or critical value of $z = \pm 1.96$ Using a critical region route: $CR < 63.2$
	dM1	Dependent on M1 A1. A correct statement – no context needed but do not allow contradicting non contextual comments. (Ignore any comparisons)
	A1	Correct conclusion in context. Must include the word claim. If they give an answer that refers to the claim then they must include the words applicants (oe), and pilots. No hypotheses then A0
		NB Award M1 A1 for a correct contextual statement on its own
ALT	B1	Both hypotheses correct in terms of p or π Must be attached to H_0 and H_1
	B1	For writing or using $N(24, 18)$ (May be implied by a correct standardisation expression)
	M1	Standardising using 28.5 or 29 or 29.5 or $x \pm 0.5$ with their mean and standard deviation (Allow \pm)
	A1	awrt 1.06 (may be applied by awrt 0.144 or 0.145) or a correct standardisation with ± 1.96 (ignore incorrect inequality symbol and allow =)
	A1	Using a probability route: awrt 0.144 or 0.145 or critical value of $z = \pm 1.96$ Using a critical region route: $CR < 32.8$
	dM1	Dependent on M1 A1. A correct statement – no context needed but do not allow contradicting non contextual comments. (Ignore any comparisons)
	A1	Correct conclusion in context. Must include the word claim. If they give an answer that refers to the claim then they must include the words applicants (oe), and pilots. No hypotheses then A0
		NB Award M1 A1 for a correct contextual statement on its own

Question Number	Scheme		Marks												
6 (a)	A sampling distribution is all the values of a statistic (obtained from a random sample) and the associated probabilities or the probability distribution of the statistic (under random sampling).		B1 (1)												
(b)	$P(6) = \frac{6}{11} \quad P(7) = \frac{3}{11} \quad P(8) = \frac{2}{11}$		B1												
	Totals (T) 12, 13, 14, 15, 16		B1												
	(6, 6) (6, 7) (6, 8) (7, 6) (7, 7) (7, 8) (8, 6) (8, 7) (8, 8)		B1												
	$[P(T = 12) =] \left(\frac{6}{11}\right)^2 = \left[\frac{36}{121}\right]$		M1												
	$[P(T = 13) =] 2 \times \left(\frac{6}{11}\right) \times \left(\frac{3}{11}\right) = \left[\frac{36}{121}\right]$														
	$[P(T = 14) =] 2 \times \left(\frac{6}{11}\right) \times \left(\frac{2}{11}\right) + \left(\frac{3}{11}\right)^2 = \left[\frac{33}{121}\right]$														
	$[P(T = 15) =] 2 \times \left(\frac{3}{11}\right) \times \left(\frac{2}{11}\right) = \left[\frac{12}{121}\right]$														
	$[P(T = 16) =] \left(\frac{2}{11}\right)^2 = \left[\frac{4}{121}\right]$														
	<table border="1"> <tr> <td>T</td> <td>12</td> <td>13</td> <td>14</td> <td>15</td> <td>16</td> </tr> <tr> <td>$P(T=t)$</td> <td>$\frac{36}{121}$</td> <td>$\frac{36}{121}$</td> <td>$\frac{33}{121}$</td> <td>$\frac{12}{121}$</td> <td>$\frac{4}{121}$</td> </tr> </table>	T	12	13	14	15	16	$P(T=t)$	$\frac{36}{121}$	$\frac{36}{121}$	$\frac{33}{121}$	$\frac{12}{121}$	$\frac{4}{121}$		A1 (7)
T	12	13	14	15	16										
$P(T=t)$	$\frac{36}{121}$	$\frac{36}{121}$	$\frac{33}{121}$	$\frac{12}{121}$	$\frac{4}{121}$										
(c)	$E(T) = "12" \times \frac{36}{121} + "13" \times \frac{36}{121} + "14" \times \frac{33}{121} + "15" \times \frac{12}{121} + "16" \times \frac{4}{121}$		M1												
	$= \frac{1606}{121} = \frac{146}{11} = 13.272\dots$		awrt 13.3 A1 (2)												
Notes			Total 10												
6 (a)	B1	A correct explanation with the words in bold													
(b)	B1	Correct probabilities – may be seen in an equation or implied by a correct probability for $T = 14$													
	B1	All 5 totals correct with no extras													
	B1	All 6 basic combinations correct, either seen or used (may be implied by correct probabilities) Allow S for 6, M for 7 and L for 8													
	M1	Correct method for one probability fit their $P(6)$, $P(7)$ and $P(8)$ If these are not stated then they must be correct													
	M1	Correct method for three of the five probabilities fit their $P(6)$, $P(7)$ and $P(8)$ If these are not stated then they must be correct													
	M1	Correct method for all five probabilities fit their $P(6)$, $P(7)$ and $P(8)$ If these are not stated then they must be correct or 5 probabilities that add up to 1													
	A1	cao Need not be in a table but probabilities must be attached to the correct total													
(c)	M1	Use of $\sum tP(T = t)$ two or more products fit their table													
	A1	awrt 13.3 (Allow $\frac{146}{11}$ oe)													

Question Number	Scheme		Marks
7 (a)	$P(L \geq 4.5) \Rightarrow P(A \geq 20.25)$		
	$P(A \geq 20.25) = (30 - 20.25) \times \frac{1}{20}$		M1
	$= 0.4875$		A1
			(2)
(b)	$\text{Var}(L) = E(L^2) - E(L)^2$		
	$[E(L^2) = E(A)] = 20$		B1
		$g(L) = \begin{cases} \frac{L}{10} & \sqrt{10} \leq L \leq \sqrt{30} \\ 0 & \text{otherwise} \end{cases}$	
	$E(L) = E(\sqrt{A}) = \frac{1}{20} \int_{10}^{30} \sqrt{a} \, dA$	$E(L) = \frac{1}{10} \int_{\sqrt{10}}^{\sqrt{30}} L^2 \, dL$	M1
	$= \frac{1}{20} \left[\frac{2}{3} a^{\frac{3}{2}} \right]_{10}^{30}$	$\frac{1}{10} \left[\frac{L^3}{3} \right]_{\sqrt{10}}^{\sqrt{30}}$	A1
	$= 4.4231\dots$		A1
	$\text{Var}(L) = "20" - ("4.4231\dots")^2$		M1
	$= 0.4358\dots$	awrt 0.436	A1
			(6)
Notes			Total 8
7 (a)	M1	$(30 - 20.25) \times \frac{1}{20}$	
	A1	cao (Allow 0.488 or $\frac{39}{80}$)	
(b)	B1	For 20	
	M1	Attempt to integrate $\frac{1}{20} \int_{10}^{30} \sqrt{a} \, dA$ or $\frac{1}{10} \int_{\sqrt{10}}^{\sqrt{30}} L^2 \, dL$ Ignore limits and accept any letter	
	A1	Fully correct integration. Accept any letter. Must have limits	
	A1	4.42 or better	
	M1	Use of $\text{Var}(L) = E(L^2) - E(L)^2$ ft their $E(L^2)$ and $E(L)$ provided $\text{Var}(L) > 0$	
	A1	awrt 0.436	