



Pearson
Edexcel

Mark Scheme (Results)

October 2023

Pearson Edexcel International Advanced Level
in Pure Mathematics (WMA11) Paper 01

Edexcel and BTEC Qualifications

Edexcel and BTEC qualifications are awarded by Pearson, the UK's largest awarding body. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information visit our qualifications websites at www.edexcel.com or www.btec.co.uk. Alternatively, you can get in touch with us using the details on our contact us page at www.edexcel.com/contactus.

Pearson: helping people progress, everywhere

Pearson aspires to be the world's leading learning company. Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: www.pearson.com/uk

October 2023

Question Paper Log Number 74316

Publications Code WMA11_01_rms_20240118

All the material in this publication is copyright

© Pearson Education Ltd 2023

General Marking Guidance

- All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.
- Mark schemes should be applied positively. Candidates must be rewarded for what they have shown they can do rather than penalised for omissions.
- Examiners should mark according to the mark scheme not according to their perception of where the grade boundaries may lie.
- There is no ceiling on achievement. All marks on the mark scheme should be used appropriately.
- All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e. if the answer matches the mark scheme. Examiners should also be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.
- Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.
- When examiners are in doubt regarding the application of the mark scheme to a candidate's response, the team leader must be consulted.
- Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.

EDEXCEL IAL MATHEMATICS

General Instructions for Marking

1. The total number of marks for the paper is 75.
2. The Edexcel Mathematics mark schemes use the following types of marks:
 - **M** marks: Method marks are awarded for 'knowing a method and attempting to apply it', unless otherwise indicated.
 - **A** marks: Accuracy marks can only be awarded if the relevant method (M) marks have been earned.
 - **B** marks are unconditional accuracy marks (independent of M marks)
 - Marks should not be subdivided.

3. Abbreviations

These are some of the traditional marking abbreviations that will appear in the mark schemes and can be used if you are using the annotation facility on ePEN.

- bod – benefit of doubt
- ft – follow through
- the symbol \checkmark will be used for correct ft
- cao – correct answer only
- cso – correct solution only. There must be no errors in this part of the question to obtain this mark
- isw – ignore subsequent working
- awrt – answers which round to
- SC: special case
- oe – or equivalent (and appropriate)
- d... or dep – dependent
- indep – independent
- dp decimal places
- sf significant figures
- * The answer is printed on the paper or ag- answer given
- \square or d... The second mark is dependent on gaining the first mark

4. All A marks are 'correct answer only' (cao.), unless shown, for example, as A1 ft to indicate that previous wrong working is to be followed through. After a misread however, the subsequent A marks affected are treated as A ft, but manifestly absurd answers should never be awarded A marks.
5. For misreading which does not alter the character of a question or materially simplify it, deduct two from any A or B marks gained, in that part of the question affected. If you are using the annotation facility on ePEN, indicate this action by 'MR' in the body of the script.
6. If a candidate makes more than one attempt at any question:
 - If all but one attempt is crossed out, mark the attempt which is NOT crossed out.
 - If either all attempts are crossed out or none are crossed out, mark all the attempts and score the highest single attempt.
7. Ignore wrong working or incorrect statements following a correct answer.

General Principles for Pure Mathematics Marking

(NB specific mark schemes may sometimes override these general principles)

Method mark for solving 3 term quadratic:

1. Factorisation

$(x^2 + bx + c) = (x + p)(x + q)$, where $|pq| = |c|$, leading to $x = \dots$

$(ax^2 + bx + c) = (mx + p)(nx + q)$, where $|pq| = |c|$ and $|mn| = |a|$, leading to $x = \dots$

2. Formula

Attempt to use correct formula (with values for a , b and c).

3. Completing the square

Solving $x^2 + bx + c = 0$: $(x \pm \frac{b}{2})^2 \pm q \pm c$, $q \neq 0$, leading to $x = \dots$

Method marks for differentiation and integration:

1. Differentiation

Power of at least one term decreased by 1. ($x^n \rightarrow x^{n-1}$)

2. Integration

Power of at least one term increased by 1. ($x^n \rightarrow x^{n+1}$)

Use of a formula

Where a method involves using a formula that has been learnt, the advice given in recent examiners' reports is that the formula should be quoted first.

Normal marking procedure is as follows:

Method mark for quoting a correct formula and attempting to use it, even if there are small mistakes in the substitution of values.

Where the formula is not quoted, the method mark can be gained by implication from correct working with values, but may be lost if there is any mistake in the working.

Exact answers

Examiners' reports have emphasised that where, for example, an exact answer is asked for, or working with surds is clearly required, marks will normally be lost if the candidate resorts to using rounded decimals.

Answers without working

The rubric says that these may not gain full credit. Individual mark schemes will give details of what happens in particular cases. General policy is that if it could be done "in your head", detailed working would not be required. Most candidates do show working, but there are occasional awkward cases and if the mark scheme does not cover this, please contact your team leader for advice.

Question Number	Scheme	Marks
1a	$y = 5x^3 + \frac{3}{x^2} - 7x = 5x^3 + 3x^{-2} - 7x$ $\left(\frac{dy}{dx} =\right) 15x^2 - 6x^{-3} - 7$	M1A1A1
		(3)
b	$\left(\frac{d^2y}{dx^2} =\right) 30x + 18x^{-4}$	M1A1
		(2)
		(5 marks)

Do not be concerned by the labelling of their answers in part (a) or part (b)

(a)

M1: Decreases the power by 1 on one of the terms $\dots x^3 \rightarrow \dots x^2$ $\dots x^{-2} \rightarrow \dots x^{-3}$ $\pm 7x \rightarrow \pm 7$
The index does not have to be processed. e.g. $\dots x^3 \rightarrow \dots x^{3-1}$

A1: Two correct unsimplified terms (indices must be processed). May be seen on different lines or listed. Condone $7x^0$ as an unsimplified term. Do not withhold this mark if there is a $+c$ present.

A1: $15x^2 - 6x^{-3} - 7$ or simplified equivalent. Condone any spurious notation around the expression but withhold this mark if there is a constant of integration. Isw once a correct answer is seen. Do not accept $15x^2 - 6x^{-3} - 7x^0$ or e.g. $15x^2 + -6x^{-3} - 7$

(b)

M1: Decreases the power by 1 on one of the terms of their changed function. Do not allow this mark to be scored for a constant of integration in (a) $\rightarrow 0$ in (b)

A1: $30x + 18x^{-4}$ or simplified equivalent e.g. $30x + \frac{18}{x^4}$. Condone e.g. $30x^1 + \frac{18}{x^4}$. Isw once a correct answer is seen. Withhold this mark if there is a constant of integration and it has not already been penalised in (a).
Allow this mark to be scored following $15x^2 - 6x^{-3} \pm 7$ in (a). Also allow full marks to be scored if they have no constant.

Note: (a) $15x^2 - 6x^{-1} - 7$ and (b) $30x + 6$ will score M1A1A0 M1A0

Question Number	Scheme	Marks
2a	$\frac{1}{8}x$	B1
		(1)
b	$\frac{1}{256}x^{\frac{3}{2}}$	B1
		(1)
c	$\left(\frac{1}{2}\left(\frac{1}{64}x^2 \times \frac{16}{\sqrt{x}}\right)\right)^{\frac{4}{3}} = \left(\frac{1}{8}x^{\frac{3}{2}}\right)^{\frac{4}{3}} = 16x^{-2}$	M1A1
		(2)
		(4 marks)

(a)

B1: $\frac{1}{8}x$ or simplified equivalent e.g. $0.125x$ or $\frac{1}{8}x^1$ Accept e.g. $\frac{x}{8}$ but $\pm\frac{1}{8}x$ is B0.

Do not withhold this mark unless it is clear that they intend to write $\frac{1}{8x}$

(b)

B1: $\frac{1}{256}x^{\frac{3}{2}}$ or simplified equivalent e.g. $0.00390625x^{\frac{3}{2}}$ condone $\frac{x^{\frac{3}{2}}}{256}$. Isw after a correct answer is seen.

Do not accept e.g. $\frac{1}{256}\sqrt{x^3}$ or $\frac{1}{256}(\sqrt{x})^3$ or $\frac{1}{256}x\sqrt{x}$ as n is not a simplified constant.

Do not accept $x^{\frac{3}{2}} \div 256$ or $\frac{1}{256x^{\frac{3}{2}}}$ (not of the required form)

Do not withhold this mark unless it is clear that they intend to write $\frac{1}{256x^{\frac{3}{2}}}$

(c)

M1: Attempts to use the index laws and proceeds to either

- $\left(\dots x^{\frac{3}{2}}\right)^{\frac{4}{3}}$ or $\left(\dots x^{-\frac{3}{2}}\right)^{\frac{4}{3}}$
- $\dots x^{\frac{8}{3}} \times \dots x^{\frac{2}{3}}$ (or $\dots x^{-\frac{8}{3}} \div \dots x^{-\frac{2}{3}}$)
- $\dots x^{-2}$
- $16x^n$ where $n \neq -\frac{1}{2}$

Allow $\frac{16}{x^2}$ to score M1.

Be aware that incorrect understanding of indices such as $\frac{16}{\sqrt{x}} = 16x^{-2}$ is M0A0

A1: $16x^{-2}$ (from a correct method)

(Correct answer with no incorrect working seen can score M1A1). Isw after a correct

answer is seen. $\frac{16}{x^2}$ is A0

Question Number	Scheme	Marks
3a	$\frac{8-\sqrt{15}}{2\sqrt{3}+\sqrt{5}} \times \frac{2\sqrt{3}-\sqrt{5}}{2\sqrt{3}-\sqrt{5}} = \frac{16\sqrt{3}-8\sqrt{5}-2\sqrt{45}+\sqrt{75}}{12-5}$	M1
	e.g. $\frac{21\sqrt{3}-14\sqrt{5}}{7}$	dM1
	$3\sqrt{3}-2\sqrt{5}$	A1
		(3)
b	$(x+5\sqrt{3})\sqrt{5} = 40-2x\sqrt{3} \Rightarrow x\sqrt{5}+2x\sqrt{3} = 40-5\sqrt{15}$	M1
	$(x =) \frac{40-5\sqrt{15}}{2\sqrt{3}+\sqrt{5}}$	A1
	$(x =) 15\sqrt{3}-10\sqrt{5}$	A1ft
		(3)
		(6 marks)

(a) Note that $3\sqrt{3}-2\sqrt{5}$ with no working scores M0dM0A0

M1: Attempts to rationalise the denominator by multiplying both numerator and denominator by $k(2\sqrt{3}-\sqrt{5})$, where

k is an integer usually 1, and proceeds to a fraction such as $\frac{\dots\sqrt{3}\pm\dots\sqrt{5}\pm\dots\sqrt{45}\pm\dots\sqrt{75}}{\dots}$ or $\frac{\dots}{12-5}$ or $\frac{\dots}{7}$

but not for e.g. $\frac{\dots}{12+2\sqrt{15}-2\sqrt{15}-5}$ or $\frac{\dots}{(2\sqrt{3})^2-(\sqrt{5})^2}$ as these still have surds on the denominator.

Allow $\sqrt{45}$ and $\sqrt{75}$ to be written in terms of $\sqrt{3}$ and $\sqrt{5}$ as well.

Condone slips in multiplying out as well as miscopying errors.

Note $\frac{21\sqrt{3}-14\sqrt{5}}{7}$ with no intermediate working can still score M1 for the denominator.

Attempting to multiply by a multiple of $\frac{2\sqrt{3}+\sqrt{5}}{2\sqrt{3}+\sqrt{5}}$ is M0dM0A0

dM1: Attempts to simplify surds and may collect terms to achieve a fraction where

- the numerator is in terms of $\sqrt{3}$ and $\sqrt{5}$ only.
- the denominator is a multiple of 7 (which may be unsimplified)

It is dependent on the previous method mark.

If they have not fully multiplied out the numerator (or implied) then this mark cannot be scored. e.g.

$$\frac{(8-\sqrt{15})(2\sqrt{3}-\sqrt{5})}{\dots} = \frac{16\sqrt{3}+8\sqrt{5}}{\dots}$$
 scores a maximum of M1dM0A0.

Note $\frac{21\sqrt{3}-14\sqrt{5}}{7}$ **with no previous working seen** scores M1dM0A0 because they have not shown any multiplying out of the brackets on the numerator.

Note $\frac{8-\sqrt{15}}{2\sqrt{3}+\sqrt{5}} = \frac{16\sqrt{3}-8\sqrt{5}-2\sqrt{45}+\sqrt{75}}{7} = 3\sqrt{3}-2\sqrt{5}$ is M1dM0A0 because they have not collected terms

on the numerator (or changed them all to be in terms of $\sqrt{3}$ and $\sqrt{5}$ before simplifying to the final answer).

A1: $3\sqrt{3} - 2\sqrt{5}$. The answer does not imply the method marks. Do not withhold this mark for slips in working such as invisible brackets provided they are recovered/implied by further work.

Note that a number of candidates are misreading $\frac{8 - \sqrt{15}}{2\sqrt{3} + 5} \times \frac{2\sqrt{3} - 5}{2\sqrt{3} - 5} = \frac{16\sqrt{3} - 40 - 2\sqrt{45} + 5\sqrt{15}}{12 - 25}$ this can score SC100 condoning one sign slip in their multiplying out of the brackets in the numerator.

(b) **Note that $x = 15\sqrt{3} - 10\sqrt{5}$ with no working scores M0A0A0**

M1: Multiplies out the brackets and isolates the x terms on one side. Condone for this mark if the surds are converted to rounded decimals.

e.g. $(\sqrt{5} + 2\sqrt{3})x = 5(8 - \sqrt{15})$ scores M1

Alternatively divides both sides by $\sqrt{5}$ and isolates the x terms on one side.

e.g. $x + 2x \frac{\sqrt{3}}{\sqrt{5}} = \frac{40}{\sqrt{5}} - 5\sqrt{3}$

Condone slips in their working and invisible brackets, but there must be two terms on each side of the equation.

A1: $x = \frac{40 - 5\sqrt{15}}{2\sqrt{3} + \sqrt{5}}$ or exact equivalent (which cannot be $15\sqrt{3} - 10\sqrt{5}$).

In the alternative method it may be seen as $x = \frac{\frac{40}{\sqrt{5}} - 5\sqrt{3}}{1 + 2 \frac{\sqrt{3}}{\sqrt{5}}}$ which can score this mark.

Do not accept proceeding from $(\sqrt{5} + 2\sqrt{3})x = 5(8 - \sqrt{15}) \Rightarrow x = 15\sqrt{3} - 10\sqrt{5}$ without any intermediate expression for x . This scores A0 as the use of calculator technology is not acceptable. An expression in decimals only is A0.

A1ft: $x = 15\sqrt{3} - 10\sqrt{5}$ only (or exact simplified equivalent) e.g. $x = 5(3\sqrt{3} - 2\sqrt{5})$ This mark cannot be awarded without the previous A mark being scored.

Follow through on their part (a) answer of the form $a\sqrt{3} + b\sqrt{5}$ so award for $5a\sqrt{3} + 5b\sqrt{5}$ or including $5(a\sqrt{3} + b\sqrt{5})$ where a and b may be fractions. Isw once they have achieved the correct answer.

Note: If a candidate has an unsimplified answer in part (a) e.g. $\frac{21\sqrt{3} - 14\sqrt{5}}{7}$ then do not withhold this mark in


part (b) if they proceed to e.g. $\frac{105\sqrt{3} - 70\sqrt{5}}{7}$ as they have already been penalised once for not giving an answer in simplest form.

Alternative method – squaring both sides (send to review if unsure)

M1: Attempts to square both sides and proceeds to a three term quadratic in x with terms collected on one side of the equation (condone slips in multiplying out and collecting terms).

A1: As above in main scheme

A1ft: As above in main scheme

Question Number	Scheme	Marks
4a	$x = -2$	B1
		(1)
b	$x^3 + 4x^2 + 4x = x(x+2)^2$	M1A1
		(2)
c		B1B1B1
		(3)
d	2 as the graphs intersect (each other) twice (since $(x+2)(x^3 + 4x^2 + 4x) = 1$ is the same as $x^3 + 4x^2 + 4x = \frac{1}{x+2}$)	B1
		(1)
		(7 marks)

Note: Check for answers by the questions or on the Figure (rather than the diagram). If there is a contradiction between the answers by the question and the main body of the solution then the main body of the solution takes precedence.

(a)

B1: $x = -2$ only. Accept labelled on the diagram but do not accept just -2 . If other equations are given (other than $y = 0$) then B0.

Beware that $x = -2$ is one of the solutions to the cubic so the equation of the asymptote must be seen in (a), on the diagram or by the question.

(b)

M1: Attempts to take out a linear factor eg $x(x^2 \pm 4x \pm 4)$ or $(x+2)(x^2 \pm 2x)$. May be implied by the correct answer.


A1: $x(x+2)^2$ or $x(x+2)(x+2)$. Accept $(x+2)(x+2)x$ and condone a missing trailing bracket on the final linear factor e.g. $x(x+2)(x+2$ but $x(x+2)^2$ is A0.

Condone $(x+0)(x+2)^2$. Isw once a correct expression is seen and ignore a spurious $= 0$

(c) **Note: If there are multiple attempts then score the highest one**

B1: Sketches a positive cubic anywhere on a set of axes. Do not penalise poor curvature provided the intention is clear and ignore if the curvature looks asymptotic. The cubic does not need to have two turning points for this mark.

Examples of acceptable cubic shapes (including where poor curvature would be condoned):



(If you are unsure then send to review)

B1: A cubic which

- has a turning point where the x -axis and vertical asymptote intersect
- passes through the origin (from quadrants 3 to 1 or from quadrants 2 to 4 – it cannot start or stop at the origin) It also cannot be a turning point.

Do not penalise poor curvature provided the intention is clear and ignore if the curvature looks asymptotic.

B1: $(-2, 0)$ **indicated on the graph** where their graph crosses or turns on the **negative x -axis** Do not be concerned regarding any other points where the graph crosses or turns on the x -axis including the origin.

Condone -2 labelled on the x -axis or the coordinates the wrong way round as $(0, -2)$ or with missing brackets.

The asymptote labelled $x = -2$ does not score this mark and do not be concerned with the point of intersection relative to the asymptote.

(d) **This mark can only be scored provided B1B1B0 or B1B1B1 is scored in (c)**


B1: **2** (real) roots **and** a valid reason e.g. (graphs/curves/functions/equations/they) **intersect/meet/cross/touch** (each other) **twice**. e.g. “intersect twice” or “2 intersections” can score this mark.

May also refer to an intersection in the first and third quadrants.

Only withhold the mark if it is clear that they are referring to intersections between the cubic graph and the x -axis.

Do not accept responses relating to the roots of the equation, use of the discriminant or other algebraic methods which do not use the graphs.

(b) Note there are a variety of different methods to finding the area of triangle ABC



B1: $\theta = \text{awrt } 42^\circ$ seen or implied. Accept 42.5. May work in radians (awrt 0.73/0.74 radians)

M1: Attempts to find AB , AD or AC using a correct method. Angles and lengths must be in the correct positions in the relevant formula or formulae. Condone slips in any rearrangement, calculations and substituting in $x = 2\sqrt{3}$ provided the method is correct. Condone working in radians provided the angles are consistently in degrees or radians within the expression or formula.

A1: awrt 6.75/6.76 or awrt 9.53/9.54 or awrt 5.07/5.08 (may be implied by further work).

dM1: Correct full method to find the total area (the expression is sufficient. It is dependent on the previous method mark. Condone use of incorrectly rounded angles/slips and may work in radians.

$$\text{e.g. Area} = \frac{1}{2} \times 5 \times "6.75" \times \sin\left(\pi - \frac{\pi}{6} - "0.74"\right)$$

May find the areas of the two separate triangles ABD and BDC and add them together.

$$\text{e.g. Area} = \frac{1}{2} \times "5.07" \times "6.75" \times \sin 30 + \frac{1}{2} \times "4.46" \times 5 \times \sin "42.5"$$

Use the diagrams above to help with the various methods. Invisible brackets may be implied by further work or their answer.

Note that if, as part of their method to find the total area, they find angle ADB but incorrectly deduce this as an acute angle then this is dM0

A1: awrt 16.1 m^2 (condone lack of units)

Question Number	Scheme	Marks
6a	$4(p-2x) = \frac{12+15p}{x+p}$ $8x^2 + 4px + 12 + 15p - 4p^2 = 0$ <p>e.g. $a=8, b=4p, c=12+15p-4p^2$</p> $(4p)^2 - 4 \times 8 \times (12+15p-4p^2) > 0$ $3p^2 - 10p - 8 > 0 \quad *$	M1
		dM1
		A1*
		(3)
b	<p>e.g. $(p-4)(3p+2) = 0 \Rightarrow 4, -\frac{2}{3}$</p> $p < -\frac{2}{3} \quad \text{or} \quad p > 4$	M1
		M1A1
		(3)
		(6 marks)

Allow part (b) to follow on from (a) without any labelling of specific parts

(a)

M1: Multiplies by $(x+p)$, multiplies out the brackets and collects terms on one side.

Condone slips and the omission of $= 0$, but there must be an $\dots x^2$ term.

May be implied by further work e.g. their attempt at the discriminant.

dM1: Attempts $b^2 - 4ac$ for their quadratic in x with their b and their c both being in terms of p . The values/expressions for a, b and c for their quadratic in x must be embedded in the correct positions in the expression but condone invisible brackets and sign slips miscopying. It is dependent on the previous method mark.

Condone for this mark if they work in terms of x instead of p provided the coefficients were actually those in terms of p from their original quadratic. i.e.

$$8x^2 + 4px + 12 + 15p - 4p^2 = 0 \Rightarrow a=8, b=4p, c=12+15p-4p^2$$

$$\Rightarrow (4x)^2 - 4 \times 8 \times (12+15x-4x^2) > 0 \quad (\text{they have just written } x \text{ instead of } p \text{ which is fine})$$

A1*: Achieves $3p^2 - 10p - 8 > 0$ with all stages of working shown:

- Multiplying out $(p-2x)(x+p)$
- Collecting terms on one side of an equation ($=0$ may be implied)
- Finding the discriminant
- At least one intermediate stage of working between the discriminant and the final answer

There must be no errors seen including invisible brackets but condone a missing trailing bracket if it does not alter the processing. e.g. $(4p)^2 - 4 \times 8 \times (12+15p-4p^2)$

Condone the > 0 appearing for the first time in their final answer, provided an incorrect inequality has not been used in earlier working.

Do not withhold this mark if they state e.g. " $b^2 - 4ac = 0$ " as part of their working.

(b)

M1: Attempts to find critical values by solving the given quadratic by either

- factorising (do not accept $(p-4)(p+\frac{2}{3})$ unless they have divided both sides of the inequality/equation by 3 first)
- completing the square $3\left(p-\frac{5}{3}\right)^2 \pm \dots \Rightarrow (p =) \dots$
- quadratic formula

Usual rules apply (see general marking principles for guidance). May be in another variable. Allow this mark to be scored if seen in (a).

This mark cannot be awarded from directly using a calculator and stating the roots.

M1: Attempts to find the outside region for their critical values. May use another variable e.g. x

May be implied by $p \leq -\frac{2}{3}$, $p \geq 4$ or incorrect use of inequalities e.g. $4 < p < -\frac{2}{3}$

This is **not** dependent on the first method mark so if a calculator has been used then this mark can still be scored.

A1: $p < -\frac{2}{3}$ or $p > 4$ (or equivalent). (may appear on separate lines). Isw once a correct

answer is seen, provided there is no contradiction and no part of the range is rejected.

This can only be scored provided both previous method marks have been awarded.

Must be in terms of p

Accept e.g. $p < -\frac{2}{3}$, $p > 4$, $p < -\frac{2}{3}$ $p > 4$ $\left\{ p : p < -\frac{2}{3} \cup p > 4 \right\}$ or variations of

these. Accept $\left(-\infty, -\frac{2}{3}\right)$, $(4, \infty)$

Do not accept e.g. $p < -\frac{2}{3}$ and $p > 4$ $\left\{ p : p < -\frac{2}{3} \cap p > 4 \right\}$ or variations of these

Do not accept $\left(-\infty, -\frac{2}{3}\right]$, $[4, \infty)$ $p \leq -\frac{2}{3}$, $p \geq 4$ $4 < p < -\frac{2}{3}$

Question Number	Scheme	Marks
7ai	$f'(4) = \frac{4(4)^2 + 10 - 7(4)^{\frac{1}{2}}}{4(4)^{\frac{1}{2}}} = \frac{15}{2}$	B1
ii	$"\frac{15}{2}" \rightarrow -\frac{2}{15}$	M1
	$y + 1 = "-\frac{2}{15}"(x - 4)$	M1
	$2x + 15y + 7 = 0$	A1
		(4)
b	$\frac{4x^2 + 10 - 7x^{\frac{1}{2}}}{4x^{\frac{1}{2}}} = \pm \dots x^{\frac{3}{2}} \pm \dots x^{-\frac{1}{2}} \pm \dots$	M1
	Two of the terms of $x^{\frac{3}{2}} + \frac{5}{2}x^{-\frac{1}{2}} - \frac{7}{4}$	A1
	$\int \left(x^{\frac{3}{2}} + \frac{5}{2}x^{-\frac{1}{2}} - \frac{7}{4} \right) dx = \frac{2}{5}x^{\frac{5}{2}} + 5x^{\frac{1}{2}} - \frac{7}{4}x (+c)$	dM1A1ft
	$\frac{2}{5}(4)^{\frac{5}{2}} + 5(4)^{\frac{1}{2}} - \frac{7}{4}(4) + c = -1 \Rightarrow c = \dots$	ddM1
	$(f(x) =) \frac{2}{5}x^{\frac{5}{2}} + 5x^{\frac{1}{2}} - \frac{7}{4}x - \frac{84}{5}$	A1
		(6)
		(10 marks)

Mark (a) and (b) together so do not be concerned with labelling of the parts

(a)

(i)

B1: $\frac{15}{2}$ oe stated (as the gradient of $f(x)$ at P).

(ii)

M1 Finds the negative reciprocal of their gradient in part (i). If they do not have a gradient in (i) then only allow $-\frac{2}{15}$ or an attempt at $-\frac{1}{f'(4)}$

M1 Attempts to find the equation of the normal using a changed gradient to that found in (i) and $(4, -1)$ with the coordinates in the correct positions. If they do not have a gradient in (i) then allow any gradient $\neq \frac{15}{2}$. If they use $y = mx + c$ they must proceed as far as $c = \dots$

A1: $2x + 15y + 7 = 0$ or any multiple of this where all the coefficients are integers and all terms are on the same side of the equation. e.g. $30y + 14 + 4x = 0$ scores A1

(b)

M1: Splits into **three** separate terms with at least one term with the correct index. The index does not need to be processed. e.g. $x^{2-\frac{1}{2}}$

A1: Two of $x^{\frac{3}{2}} + \frac{5}{2}x^{-\frac{1}{2}} - \frac{7}{4}$ (unsimplified but the indices must be processed). May appear as a list of terms on different lines. May be implied by a correctly integrated expression.

dM1: Attempts to increase the power by one on at least one term. It is dependent on the previous method mark. The index does not need to be processed for this mark. It cannot be scored for attempting to integrate individual terms on the numerator or denominator.

A1ft: $\frac{2}{5}x^{\frac{5}{2}} + 5x^{\frac{1}{2}} - \frac{7}{4}x$ or exact unsimplified equivalent (indices processed). e.g. $\frac{x^{\frac{5}{2}}}{\frac{5}{2}} + \frac{\frac{5}{2}x^{\frac{1}{2}}}{\frac{1}{2}} - \frac{7}{4}x$

Follow through on an expression of the form $Ax^{\frac{3}{2}} + Bx^{-\frac{1}{2}} + C \rightarrow \frac{Ax^{\frac{5}{2}}}{\frac{5}{2}} + \frac{Bx^{\frac{1}{2}}}{\frac{1}{2}} + Cx$ where A ,

B and C are all non-zero.

Do not be concerned with the presence or omission of the constant of integration.

Accept x^1 for x .

Ignore any spurious notation including the presence of the integral sign.

ddM1: Attempts to substitute $x=4$ into their changed expression (condone one slip in substituting in), sets equal to -1 and attempts to find c .

This mark cannot be scored if they do not have a constant of integration.

It is dependent on both of the previous method marks.

Do not be concerned by the mechanics of the rearrangement and condone arithmetical slips, but they must achieve a value.

The substitution may be implied by a correct value for c for their integrated expression if no working is shown (you may need to check this), or allow for $\pm \frac{84}{5}$

A1: $(f(x) =) \frac{2}{5}x^{\frac{5}{2}} + 5x^{\frac{1}{2}} - \frac{7}{4}x - \frac{84}{5}$ or any equivalent expression e.g. $0.4x^{\frac{5}{2}} + 5x^{\frac{1}{2}} - 1.75x - 16.8$

Accept x^1 for x . Condone equivalent fractions for the coefficients provided both numerator

and denominator are integers. e.g. $(f(x) =) \frac{x^{\frac{5}{2}}}{2.5} + 5x^{\frac{1}{2}} - \frac{7}{4}x - \frac{84}{5}$ is A0

Withhold the final mark if there is still integration notation around the answer or if it is set equal to 0

Question Number	Scheme	Marks
8a	$x\left(x^3 - \frac{7}{2}x - 5\right) = \frac{15}{2} - 5x$ $\Rightarrow 2x^4 - 7x^2 - 15 = 0 \quad *$	M1 A1*
		(2)
b	$(2x^2 + 3)(x^2 - 5) = 0 \Rightarrow (x^2 =) \dots$ $(x =) (\pm)\sqrt{5}$	M1 B1
	$y = \frac{15}{2 \times (\pm)\sqrt{5}} - 5 = \dots \quad \text{or} \quad y = \left(\pm\sqrt{5}\right)^3 - \frac{7}{2}\left(\pm\sqrt{5}\right) - 5$	M1
	$\left(-\sqrt{5}, -\frac{3}{2}\sqrt{5} - 5\right) \quad \left(\sqrt{5}, \frac{3}{2}\sqrt{5} - 5\right)$ $PQ = \sqrt{("2\sqrt{5}")^2 + ("3\sqrt{5}")^2} = \dots$	dM1
	$PQ = \sqrt{65}$	A1cao
		(5)
		(7 marks)

(a)

M1: Sets up a correct equation in x . Invisible brackets may be implied by further work.

e.g. $x^4 - \frac{7}{2}x^2 - 5x = \frac{15}{2} - 5x$

A1*: Multiplies out and rearranges the equation to achieve the given answer with no errors seen including invisible brackets. Condone the implied $= 0$ in their working, provided it is present in their final answer. There must be at least one stage of intermediate working between their starting equation and proceeding to the given answer.

$x^4 - \frac{7}{2}x^2 - 5x = \frac{15}{2} - 5x \Rightarrow 2x^4 - 7x^2 - 15 = 0$ is M1A0*

(b) **Work for (b) may be seen in (a) which can score**

M1: Attempts to solve the quadratic equation in x^2 by either

- factorising
- completing the square
- quadratic formula

Usual rules apply. Do not be too concerned by the labelling e.g. it is acceptable to use a different variable as well as condoning $y = x^2$ and $x = x^2$

This mark cannot be awarded from directly using a calculator and stating the roots.

B1: At least one of $\sqrt{5}$ and $-\sqrt{5}$ (and no others)

M1: **Note that if they do not square root their solution to the quadratic in x^2 then this mark cannot be scored (and no further marks).**

Attempts to substitute in $\pm\sqrt{"5"}$ (which must be of the form $+k\sqrt{b}$ and $-k\sqrt{b}$ where k may be 1) into either of the equations of the curves and proceeds to find irrational expressions for the y coordinates of P and Q of the form $\pm 5 \pm n\sqrt{"5"}$ (which may be the same for both). They cannot be decimals.

May be seen as e.g. $\left(-\sqrt{5}, \frac{-3\sqrt{5}-10}{2}\right) \quad \left(\sqrt{5}, \frac{3\sqrt{5}-10}{2}\right)$

dM1: A full attempt to find the length PQ . It is dependent on the previous method mark.

- They must be using $(k\sqrt{b}, \pm 5 \pm n\sqrt{b})$, $(-k\sqrt{b}, \pm 5 \pm n\sqrt{b})$ or equivalent.
- Look for an expression for PQ of the form e.g.

$$\sqrt{\left((k\sqrt{b} - -k\sqrt{b})\right)^2 + \left((\pm 5 \pm n\sqrt{b}) - (\pm 5 \pm n\sqrt{b})\right)^2}$$

- They must be attempting to find $(x_2 - x_1)^2$ for their x coordinates which would lead to $\pm 2k\sqrt{b}$ but condone ambiguity in the location of negatives when calculating $(y_2 - y_1)^2$.

e.g. $x = \pm\sqrt{5}$: $\sqrt{\left((\sqrt{5} - -\sqrt{5})\right)^2 + \left(-\frac{10+3\sqrt{5}}{2} - \frac{-10+3\sqrt{5}}{2}\right)^2}$ scores B1M1dM1

e.g. $x = \pm\sqrt{11}$: $\sqrt{\left((-\sqrt{11} - \sqrt{11})\right)^2 + \left(\frac{110+15\sqrt{11}}{22} - \frac{-110-15\sqrt{11}}{22}\right)^2}$ scores B0M1dM1

(as their $(x_2 - x_1)^2$ is correct for their $\pm\sqrt{5}$ and we condone sign errors when finding $(y_2 - y_1)^2$ using their y coordinates of P and Q)

- If the subtractions are not shown then they must achieve either of the following:
 - $\sqrt{(2k\sqrt{b})^2 + (2n\sqrt{b})^2}$ (n does not have to be an integer) or equivalent
 - $\sqrt{(2k\sqrt{b})^2 + 0^2}$ (where the y coordinates were the same)
(They must have still found the y coordinates to score this mark)

Do not condone invisible brackets unless it does not affect the processing. This mark may be implied by their answer.

A1: $\sqrt{65}$ cao (**this cannot be scored if the first method mark has not been scored**)

Question Number	Scheme	Marks
9a	$0.72 = OP \times 0.6 \Rightarrow OP = 1.2$ or $\frac{0.72}{0.6} = 1.2$ or $0.72 = 1.2 \times 0.6 \Rightarrow OP = 1.2$ *	B1*
		(1)
b	$\frac{1}{2} \times (x+1.2)^2 \times 0.6$	B1
	$\frac{1}{2} \times (x+1.2)^2 \times 0.6 - \frac{1}{2} \times 1.2^2 \times 0.6 = 90 \Rightarrow 5x^2 + 12x - 1500 = 0$ *	M1A1*
c	$x = \text{awrt } 16.2 \left(= \frac{-6 + 4\sqrt{471}}{5} \right)$	B1
	Perimeter = $0.72 + "16.2" + "16.2" + 0.6 \times ("16.2" + 1.2) = \text{awrt } 43 / \text{awrt } 44$ m (see notes)	M1A1
		(3)
		(7 marks)

(a)

B1*: Either

- forms an equation using $l = r\theta$ with $l = 0.72$ and $\theta = 0.6$ and concludes $OP = 1.2$
e.g. $0.72 = r \times 0.6 \Rightarrow r = 1.2$
- states $\frac{0.72}{0.6} = 1.2$
- forms the equation using $0.72 = 1.2 \times 0.6$ and concludes that $OP = 1.2$

Condone the use of r or any other variable instead of OP . Condone lack of units or incorrect units.

Allow the angle to be in degrees instead or attempts to find the diameter first, but they must not round the value at any point in their working.

e.g. $0.72 = \frac{0.6}{360} \times 180 \pi d \Rightarrow d = 2.4 \Rightarrow r = 1.2$ scores B1 (exact angle used)

e.g. $\frac{0.72}{\left(\frac{34.4}{360}\right)} = 2\pi r \Rightarrow r = 1.2$ scores B0 (34.4 is a rounded angle)

(b)

B1: Correct expression for the area of sector $OPQRSO$

$\frac{1}{2} \times (x+1.2)^2 \times 0.6$ Missing brackets may be implied by further work.

May appear within an equation e.g. $\frac{1}{2} \times 0.6 [(x+1.2)^2 - 1.2^2] = 90$

Condone incorrect equations i.e. $\frac{1}{2} \times (x+1.2)^2 \times 0.6 = 90$

M1: Proceeds from an equation of the form $...(x \pm 1.2)^2 \times 0.6 - ... \times 1.2^2 \times 0.6 = 90$ or equivalent, (i.e. both should be positive but may be missing e.g. $\frac{1}{2}$) and attempts to multiply out the brackets. Condone slips and invisible brackets.

A1*: Achieves the given answer with no errors seen including invisible brackets. They must have multiplied out any brackets before proceeding to the given answer to score this mark.

e.g. $\frac{54}{125} + \frac{18}{25}x + \frac{3}{10}x^2 - 0.432 = 90 \Rightarrow 5x^2 + 12x - 1500 = 0$ is acceptable as a final step to score A1*

(c)

B1: $x = \text{awrt } 16.2 \left(= \frac{-6 + 4\sqrt{471}}{5} \right)$ (ignore the negative root) May be seen on the diagram or next to the quadratic in the question. It may also be implied in their working to find the perimeter. It cannot be for the values embedded in the quadratic formula.

M1: A correct method to find the perimeter using their value for x which must be positive. They must proceed as far as a value for the perimeter to score this mark. Do not condone the lengths 0.72 and 1.2 to be rounded, but condone for example their "16.2" to be "16".

May see exact values used in the expression e.g. $2 \left(\frac{-6 + 4\sqrt{471}}{5} \right) + 0.72 + \left(\frac{12\sqrt{471}}{25} \right) = \dots$

A1: awrt 43 m or awrt 44 m, **including units, following a correct method. Typically look for 43.4...m or 43.5...m**

Note some incorrect methods lead to awrt 43 which is maximum B1M0A0.

e.g. "16.2"×0.6+"16.2"+"16.2"+0.72 = 42.7 which is B1M0A0 (uses 16.2 instead of 16.2+1.2)

e.g. "16.2+1.2"×0.6+"16.2"+"16.2" = 42.8 which is B1M0A0 (missing arc *PS*)

Question Number	Scheme	Marks
10ai	$(n =) 3$	B1
ii	1080	B1
		(2)
b	$(1620, -3)$	B1B1
		(2)
c	e.g. $k = \frac{1}{2} \left(\frac{12}{5} - \frac{3}{5} \right) = \frac{9}{10}$	M1A1
		(2)
		(6 marks)

Note: answers may appear next to questions or on the diagram.

If there is a contradiction between the answer by the question/on the diagram and the main body of the work then the answer in the main body of the work takes precedence.

(a)

(i)

B1: $(n =) 3$

(ii)

B1: 1080 (must be in degrees). Do not accept e.g. (1080, 3)

(b)

B1: One of the two coordinates $(1620, -3)$ Condone missing brackets. Allow $x = \dots, y = \dots$

Condone 9π instead of 1620 for this mark and condone the coordinates to be the wrong way round e.g. $(-3, 1620)$ scores B1B0

B1: Both coordinates $(1620, -3)$ Condone missing brackets. Allow $x = \dots, y = \dots$

(c)

M1: Attempts to find a value for k by either

- attempting to find $k = \frac{1}{2} \left(\frac{12}{5} - \frac{3}{5} \right) = \dots$

- attempting to find $k = \frac{12}{5} - \frac{1}{2} \left(\frac{12}{5} + \frac{3}{5} \right) = \dots$ or $k = -\frac{3}{5} + \frac{1}{2} \left(\frac{12}{5} + \frac{3}{5} \right) = \dots$

- forming two **correct** simultaneous equations and eliminating $\sin a$ (may be labelled as a different variable, condone a for $\sin a$)

$$2 \sin(a) + k = \frac{12}{5}, \quad -2 \sin(a) + k = -\frac{3}{5} \Rightarrow k = \dots \text{ scores M1}$$

They must have correctly used $\sin(-a) = -\sin(a)$. If they eliminate k instead, they must proceed to find a or $\sin a$ and then substitute back into one of the equations to find a value for k

$$\text{i.e. } 4 \sin a = 3 \Rightarrow a = 48.59 \dots \Rightarrow 2 \sin(48.59 \dots) + k = \frac{12}{5} \Rightarrow k = \dots$$

A1: $\frac{9}{10}$ or 0.9 (this is not awrt 0.9 so if they achieve a rounded value e.g. 0.899... and round to 0.9 then M1A0)

Question Number	Scheme	Marks
11a	$2(x \pm \dots)^2$	B1
	$\dots(x \pm 3)^2 \dots$	M1
	$2(x-3)^2 - 4$	A1
		(3)
b	$(3, -4)$	B1ft
		(1)
c	$m = \frac{28 - -4}{-1 - 3} (= -8)$	M1
	$y - 28 = -8(x+1)$	dM1
	$y = -8x + 20$	A1
		(3)
d	$y \leq -8x + 20$ and $y \geq 2x^2 - 12x + 14$ (or $y \geq 2(x-3)^2 - 4$)	B1ftB1ft
	$y \leq -8x + 20$ $y \geq 2x^2 - 12x + 14$ $y \geq 0, x \geq 0$	B1cso
		(3)
		(10 marks)

Note: answers may appear next to questions or on the diagram.

If there is a contradiction between the answer by the question/on the diagram and the main body of the work then the answer in the main body of the work takes precedence.

(a)

B1: $a = 2$ which may be stated or seen in the expression $2(x \pm \dots)^2$.

M1: $\dots(x \pm 3)^2 \dots$ may be implied by $b = \pm 3$. Accept unsimplified versions e.g. $\dots \left(x \pm \frac{12}{4}\right)^2 \dots$

A1: $2(x-3)^2 - 4$ isw following a correct expression (e.g. if they proceed to state incorrect values for a, b and/or c). If they just state the values then withhold this mark. Constants must be integers.

SC110 $2(x-3) - 4$ (slip writing missing the squared on the bracket)

(b)

B1ft: $(3, -4)$ or e.g. $x = 3, y = -4$. Condone invisible brackets. Follow through from their part (a) so score for $(-b, c)$

(c)

M1: Attempts to find the gradient between their minimum point and $(-1, 28)$. Score for the expression $\frac{28 - (-4)}{-1 - (-3)}$ or equivalent. They must be subtracting the correct way round. Do not condone sign slips for this mark. If a value appears with no method (nor implied using their minimum point and $(-1, 28)$), then this mark cannot be scored (M0dM0A0)

May be part of the equation for l e.g. $\frac{28 - (-4)}{y - 28} = \frac{-1 - (-3)}{x + 1}$ or $\frac{28 - (-4)}{y + 4} = \frac{-1 - (-3)}{x - 3}$

dM1: Attempts to find the equation using their gradient. It is dependent on the previous method mark. Score for $y - 28 = -8(x + 1)$ or $y + 4 = -8(x - 3)$ but condone one sign slip substituting in the coordinates for x and y .

If they use $y = mx + c$ they must proceed as far as $c = \dots$

A1: $y = -8x + 20$

(d) Allow consistent use of either $\leq \geq$ or $< >$ Ignore use of and/or or equivalent notation for this question.

Note that $2x^2 - 12x + 14$ may be expressed as $2(x - 3)^2 - 4$

Allow the use of $f(x)$ instead of y but the use of R instead of y is not acceptable.

B1ft: One of

- $y \leq -8x + 20$ follow through on their part (c)
- $y \geq 2x^2 - 12x + 14$ follow through on their part (b) so $2(x - 3)^2 - 4$

May be part of an inconsistent set of inequalities e.g. $2x^2 - 12x + 14 < y \leq -8x + 20$

B1ft: Both

- $y \leq -8x + 20$ follow through on their part (c)
- $y \geq 2x^2 - 12x + 14$ follow through on their part (b) so $2(x - 3)^2 - 4$

May also be seen as $2x^2 - 12x + 14 \leq y \leq -8x + 20$

B1cso: $y \leq -8x + 20$ $y \geq 2x^2 - 12x + 14$ $y \geq 0, x \geq 0$ Do not isw for this mark.

Also accept $2x^2 - 12x + 14 \leq y \leq -8x + 20$ $y \geq 0, x \geq 0$

Accept inequalities $0 \leq x \leq p$ provided $p \geq \frac{5}{2}$

$0 \leq y \leq q$ provided $q \geq 20$

Note: The equations may be written in alternative forms e.g. $8x + y \leq 20$ which is also acceptable

