Please check the examination details	below l	before ente	ring your candidate information
Candidate surname			Other names
Pearson Edexcel International Advanced Level	Centre	Number	Candidate Number
Monday 21 Ja	nu	ary	2019
Afternoon (Time: 1 hour 30 minute	s)	Paper Re	eference WME01/01
Mechanics M1 Advanced/Advanced Su	bsid	iary	
You must have: Mathematical Formulae and Statis	tical Ta	ables (Blu	ue)

Candidates may use any calculator allowed by the regulations of the Joint Council for Qualifications. Calculators must not have the facility for symbolic algebra manipulation, differentiation and integration, or have retrievable mathematical formulae stored in them.

Instructions

- Use **black** ink or ball-point pen.
- If pencil is used for diagrams/sketches/graphs it must be dark (HB or B). Coloured pencils and highlighter pens must not be used.
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer **all** questions and ensure that your answers to parts of questions are clearly labelled.
- Answer the questions in the spaces provided
 there may be more space than you need.
- You should show sufficient working to make your methods clear. Answers without working may not gain full credit.
- Whenever a numerical value of g is required, take $g = 9.8 \text{ m s}^{-2}$, and give your answer to either two significant figures or three significant figures.
- When a calculator is used, the answer should be given to an appropriate degree of accuracy.

Information

- The total mark for this paper is 75.
- The marks for each question are shown in brackets
 use this as a quide as to how much time to spend on each question.

Advice

- Read each question carefully before you start to answer it.
- Try to answer every question.
- Check your answers if you have time at the end.

Turn over ▶



P54829A
©2019 Pearson Education Ltd.



1.	Two particles, A and B , have masses $2m$ and $3m$ respectively. They are moving towards each other in opposite directions along the same straight line on a smooth horizontal plane when they collide directly. Immediately before they collide, the speed of A is $3u$ and the speed of B is u . As a result of the collision, the speed of A is halved and the direction of motion of each particle is reversed.
	(i) Find the speed of B immediately after the collision.
	(ii) Find the magnitude of the impulse exerted on A by B in the collision. (6)



uestion 1 continued	
	(Total 6 marks)



2. [In this question i and j are horizontal unit vectors due east and due north respectively position vectors are given relative to a fixed origin O.]	and
At time $t = 0$, a bird A leaves its nest, that is located at the point with position ver $(20\mathbf{i} - 17\mathbf{j})$ m, and flies with constant velocity $(-6\mathbf{i} + 7\mathbf{j})$ m s ⁻¹ . At the same time a sec bird B leaves its nest which is located at the point with position vector $(-8\mathbf{i} + 9\mathbf{j})$ m flies with constant velocity $(p\mathbf{i} + 2p\mathbf{j})$ m s ⁻¹ , where p is a constant. At time $t = 4$ s, bird south west of bird A.	ond and
(a) Find the direction of motion of A , giving your answer as a bearing to the nearest deg	gree. (3)
(b) Find the speed of <i>B</i> .	(10)

	Leave
	blank
Question 2 continued	
Yassasa - sommasa	



Question 2 continued		

Question 2 continued	blank
	Q2
(Total 13 marks)	



3.




Figure 1

A lift of mass $M \, \mathrm{kg}$ is being raised by a vertical cable attached to the top of the lift. A person of mass $m \, \mathrm{kg}$ stands on the floor inside the lift, as shown in Figure 1. The lift ascends vertically with constant acceleration $1.4 \, \mathrm{m \, s^{-2}}$. The tension in the cable is $2800 \, \mathrm{N}$ and the person experiences a constant normal reaction of magnitude $560 \, \mathrm{N}$ from the floor of the lift. The cable is modelled as being light and inextensible, the person is modelled as a particle and air resistance is negligible.

(a) Write down an equation of motion for the person only.

(2)

(b) Write down an equation of motion for the lift only.

(2)

- (c) Hence, or otherwise, find
 - (i) the value of m,
 - (ii) the value of M.

(3)





Question 3 continued		Leave
Question & continued		
		Q3
	(Total 7 marks)	



4.




Figure 2

A boy sees a box on the end Q of a plank PQ which overhangs a swimming pool. The plank has mass $30\,\mathrm{kg}$, is $5\,\mathrm{m}$ long and rests in a horizontal position on two bricks. The bricks are modelled as smooth supports, one acting on the rod at P and one acting on the rod at R, where $PR = 3\,\mathrm{m}$. The support at R is on the edge of the swimming pool, as shown in Figure 2. The boy has mass $40\,\mathrm{kg}$ and the box has mass $2.5\,\mathrm{kg}$. The plank is modelled as a uniform rod and the boy and the box are modelled as particles.

The boy steps on to the plank at P and begins to walk slowly along the plank towards the box.

(a) Find the distance he can walk along the plank from P before the plank starts to tilt.

(4)

(b) State how you have used, in your working, the fact that the box is modelled as a particle.

(1)

A rock of mass M kg is placed on the plank at P. The boy is then able to walk slowly along the plank to the box at the end Q without the plank tilting. The rock is modelled as a particle.

(c) Find the smallest possible value of M.

(4)

	Leave
	blank
Question 4 continued	



Question 4 continued

Question 4 continued	Leave blank
	Q4
(Total 9 marks)	



5.




Figure 3

A small metal box of mass 6 kg is attached at B to two ropes BP and BQ. The fixed points P and Q are on a horizontal ceiling and PQ = 3.5 m. The box hangs in equilibrium at a vertical distance of 2 m below the line PQ, with the ropes in a vertical plane and with angle $BQP = 45^{\circ}$, as shown in Figure 3. The box is modelled as a particle and the ropes are modelled as light inextensible strings. Find

(i) the tension in *BP*,

(.1			DO
(11)	the	tension	1n	R()
(/	ULIU	COMBIGH		2

(10)	
------	--

	Question 5 continued	blank
Q		
Q		
Q:		
Q:		
Q		
Q		
Q		
Q		
Q		
Q		
Q		
Q		
Q		
Q		
Q		
Qé		
Q4		
Q4		
Q3		
Q4		
Q5		
Q5		
QS		
Q\$		
		Q5
(Total 10 marks)	(Total 10 marks)	



- 6. A train travels for a total of 270 s along a straight horizontal track between two stations A and B. The train starts from rest at A and moves with constant acceleration for 60 s until it reaches a speed of $V \, \text{m s}^{-1}$. The train then travels at this constant speed $V \, \text{m s}^{-1}$ before it moves with constant deceleration for 30 s, coming to rest at B.
 - (a) Sketch below a speed-time graph for the journey of the train between the two stations *A* and *B*.

(2)

Given that the distance between the two stations is 4.5 km,

(b) find the value of V,

(3)

(c) find how long it takes the train to travel from station A to the point that is exactly halfway between the two stations.

(4)

The train is travelling at speed $\frac{1}{4}V$ m s⁻¹ at times T_1 seconds and T_2 seconds after leaving station A.

(d) Find the value of T_1 and the value of T_2

(5)



	Leave blank
Question 6 continued	Dialik
4 monton o continuou	



Question 6 continued		
guestion o continued		

Question 6 continued	blank
	Q6
(Total 14 marks)	Í
(Total 14 Illal KS)	



(1)

7.




Figure 4

Two particles A and B have masses m and 3m respectively. The particles are attached to the ends of a light inextensible string. Particle A is held at rest on a rough horizontal table.

The coefficient of friction between particle A and the table is $\frac{1}{5}$. The string lies along the table and passes over a small smooth light pulley that is fixed at the edge of the table. Particle B is at rest on a rough plane that is inclined to the horizontal at an angle α , where $\tan \alpha = \frac{4}{3}$, as shown in Figure 4. The coefficient of friction between particle B and the inclined plane is $\frac{1}{3}$. The string lies in the vertical plane that contains the pulley and a line of greatest slope of the inclined plane. The system is released from rest with the string taut and B slides down the inclined plane. Given that A does not reach the pulley,

- (a) find the tension in the string, (11)
- (b) state where in your working you have used the fact that the string is modelled as being light,
- (c) find the magnitude of the force exerted on the pulley by the string. (4)



	blank
Question 7 continued	Ulalik
Question / continued	



Question 7 continued		



	Leave
	blank
Question 7 continued	



(Total 16 marks) TOTAL FOR PAPER: 75 MARKS	Question 7 continued		Leave
(Total 16 marks) TOTAL FOR PAPER: 75 MARKS	-		
(Total 16 marks) TOTAL FOR PAPER: 75 MARKS			
(Total 16 marks) TOTAL FOR PAPER: 75 MARKS			
(Total 16 marks) TOTAL FOR PAPER: 75 MARKS			
(Total 16 marks) TOTAL FOR PAPER: 75 MARKS			
(Total 16 marks) TOTAL FOR PAPER: 75 MARKS			
(Total 16 marks) TOTAL FOR PAPER: 75 MARKS			
(Total 16 marks) TOTAL FOR PAPER: 75 MARKS			
(Total 16 marks) TOTAL FOR PAPER: 75 MARKS			
(Total 16 marks) TOTAL FOR PAPER: 75 MARKS			
(Total 16 marks) TOTAL FOR PAPER: 75 MARKS			
(Total 16 marks) TOTAL FOR PAPER: 75 MARKS			
(Total 16 marks) TOTAL FOR PAPER: 75 MARKS			
(Total 16 marks) TOTAL FOR PAPER: 75 MARKS			
(Total 16 marks) TOTAL FOR PAPER: 75 MARKS			
(Total 16 marks) TOTAL FOR PAPER: 75 MARKS			
(Total 16 marks) TOTAL FOR PAPER: 75 MARKS			
(Total 16 marks) TOTAL FOR PAPER: 75 MARKS			
(Total 16 marks) TOTAL FOR PAPER: 75 MARKS			
(Total 16 marks) TOTAL FOR PAPER: 75 MARKS			
(Total 16 marks) TOTAL FOR PAPER: 75 MARKS			
(Total 16 marks) TOTAL FOR PAPER: 75 MARKS			
(Total 16 marks) TOTAL FOR PAPER: 75 MARKS			
(Total 16 marks) TOTAL FOR PAPER: 75 MARKS			
(Total 16 marks) TOTAL FOR PAPER: 75 MARKS			
TOTAL FOR PAPER: 75 MARKS			Q'
		TOTAL FOR PAPER: 75 MARKS	