



Pearson

Mark Scheme (Results)

Summer 2017

Pearson Edexcel International GCSE
In Mathematics B (4MB0) Paper 01R

Edexcel and BTEC Qualifications

Edexcel and BTEC qualifications are awarded by Pearson, the UK's largest awarding body. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information visit our qualifications websites at www.edexcel.com or www.btec.co.uk. Alternatively, you can get in touch with us using the details on our contact us page at www.edexcel.com/contactus.

Pearson: helping people progress, everywhere

Pearson aspires to be the world's leading learning company. Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: www.pearson.com/uk

Summer 2017

Publications Code 4MB0_01R_1706_MS

All the material in this publication is copyright

© Pearson Education Ltd 2017

General Marking Guidance

- All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.
- Mark schemes should be applied positively. Candidates must be rewarded for what they have shown they can do rather than penalised for omissions.
- Examiners should mark according to the mark scheme not according to their perception of where the grade boundaries may lie.
- There is no ceiling on achievement. All marks on the mark scheme should be used appropriately.
- All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e. if the answer matches the mark scheme. Examiners should also be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.
- Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.
- When examiners are in doubt regarding the application of the mark scheme to a candidate's response, the team leader must be consulted.
- Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.
- **Types of mark**
 - M marks: method marks
 - A marks: accuracy marks
 - B marks: unconditional accuracy marks (independent of M marks)
- **Abbreviations**
 - cao – correct answer only
 - ft – follow through
 - isw – ignore subsequent working
 - SC - special case
 - oe – or equivalent (and appropriate)
 - dep – dependent
 - indep – independent
 - awrt – answer which rounds to
 - eeoo – each error or omission

- **No working**

If no working is shown then correct answers normally score full marks

If no working is shown then incorrect (even though nearly correct) answers score no marks.

- **With working**

If there is a wrong answer indicated on the answer line always check the working in the body of the script (and on any diagrams), and award any marks appropriate from the mark scheme.

If it is clear from the working that the "correct" answer has been obtained from incorrect working, award 0 marks.

Any case of suspected misread loses A (and B) marks on that part, but can gain the M marks.

If working is crossed out and still legible, then it should be given any appropriate marks, as long as it has not been replaced by alternative work.

If there is a choice of methods shown, then no marks should be awarded, unless the answer on the answer line makes clear the method that has been used.

If there is no answer on the answer line then check the working for an obvious answer.

- **Ignoring subsequent work**

It is appropriate to ignore subsequent work when the additional work does not change the answer in a way that is inappropriate for the question: eg. Incorrect cancelling of a fraction that would otherwise be correct.

It is not appropriate to ignore subsequent work when the additional work essentially makes the answer incorrect eg algebra.

Transcription errors occur when candidates present a correct answer in working, and write it incorrectly on the answer line; mark the correct answer.

- **Parts of questions**

Unless allowed by the mark scheme, the marks allocated to one part of the question CANNOT be awarded in another.

Q	Working	Answer	Mark	Notes
1	(a)	57900	1	B1
	(b)	0.05	1	B1
				Total 2 marks
2	3200000 or 32×10^5		2	M1
		3.2×10^6		A1
				Total 2 marks
3	$(3x =)7 - 2$		2	M1
		$\frac{5}{3}$ (oe)		A1 $1\frac{2}{3}$, 1.6̇, 1.6̄, awrt 1.67 NB do not allow 1.66
				Total 2 marks
4	$\frac{14}{30} \times 360$ (oe)		2	M1 full method required
		168		A1
				Total 2 marks
5	$\pm \frac{96-90}{96}$ or $\pm \frac{6}{96}$ or $\frac{90}{96} \times 100$		3	M1
	$\pm \frac{96-90}{96} \times 100$ or $100 - \frac{90}{96} \times 100$			M1 DEP
		$6\frac{1}{4}$ (oe)		A1 $6.25, \frac{25}{4}$
NB -6.25 score M1M1A0				Total 3 marks

6	(a) (i)		$x \geq -5$	2	B1
	(ii)		$x < 6$		B1
	(b)	Single Line -5		1	B1 FT their inequalities allow $-5 [\text{-----}) 6$
					Total 3 marks

7		$\Delta = 5 \times -2 - 4 \times -3 (= 2)$		3	M1
		$\frac{1}{\Delta} \begin{pmatrix} -2 & 3 \\ -4 & 5 \end{pmatrix}$			M1 (Indep) allow $\Delta=1$
			$\begin{pmatrix} -1 & 1\frac{1}{2} \\ -2 & 2\frac{1}{2} \end{pmatrix} \text{(oe)}$		A1 Accept $\frac{1}{2} \begin{pmatrix} -2 & 3 \\ -4 & 5 \end{pmatrix}$
					Total 3 marks



8		$2\sqrt{3} - 2\sqrt{2} + \sqrt{6} \times \sqrt{3} - \sqrt{6} \times \sqrt{2}$ or $2\sqrt{3} - 2\sqrt{2} + \sqrt{18} - \sqrt{12}$		3	M1 Correctly expand brackets. Allow one sign error.
		$\sqrt{18} = 3\sqrt{2}$ and $\sqrt{12} = 2\sqrt{3}$ or $\sqrt{6} \times \sqrt{3} = 3\sqrt{2}$ and $\sqrt{6} \times \sqrt{2} = 2\sqrt{3}$			M1 May be embedded within working (DEP)
		$2\sqrt{3} - 2\sqrt{2} + 3\sqrt{2} - 2\sqrt{3}$	$\sqrt{2}$		A1 dep on M2 cso
NB Answers derived from decimal expansions score M0M0A0					Total 3 marks

9	tangent drawn at (2, 2)		3	M1	Must only touch curve at (2,2)
	$\frac{y \text{ increment}}{x \text{ increment}}$			M1(DEP)	Attempt to find gradient for their tangent.
		3		A1	dep on tangent drawn ft their tangent providing $2.5 \leq \text{gradient} \leq 4$
Total 3 marks					

10	Shaded area = $(2r)^2 - \pi r^2$, $0.858 r^2$ or better. OR $\frac{\pi(r^2)}{4(r^2)} \times 100$		3	M1	
	$\frac{4r^2 - \pi r^2}{4r^2} \times 100\%$ OR $100 - \frac{\pi(r^2)}{4(r^2)} \times 100$			M1 DEP	
		21.5		A1	awrt 21.4 or 21.5
Total 3 marks					

11	(a)		Q, Z, N correctly labelled	1	B1
	(b)		$\{4, \sqrt{2}, -3, \frac{2}{3}\}$ four elements in correct position	2	B2
Total 3 marks					

12		$\begin{pmatrix} 10 & 9 \\ 13 & 9 \\ -4 & -10 \end{pmatrix}$	3	M1 Correct dimension and at least one correct entry. A1 Any three correct entries. A1 All correct.
NB for a transcription error allow at most one value changed or two values transposed.			Total 3 marks	

13 (a)			1	B1 Correct region shaded.
(b)			1	B1 Correct region shaded.
(c)		$(F \cup G) \cap H'$ (oe)	1	B1 Accept $(F \cap H') \cup (G \cap H')$ or $(F \cap G' \cap H') \cup (F \cap G \cap H') \cup (F' \cap G \cap H')$ Do not accept $F \cup G \cap H'$
			Total 3 marks	

14	$\frac{6}{10} \times \frac{4}{10}$ or $\frac{4}{10} \times \frac{6}{9}$ or $\frac{6}{10} \times \frac{6}{10} + \frac{4}{10} \times \frac{3}{9}$		3	M1
	$\frac{6}{10} \times \frac{4}{10} + \frac{4}{10} \times \frac{6}{9}$ or $1 - \left(\frac{6}{10} \times \frac{6}{10} + \frac{4}{10} \times \frac{3}{9} \right)$			M1 DEP
		$\frac{38}{75}$ (oe)		A1 AWRT 0.507
				Total 3 marks

15	SF = $\sqrt[3]{\frac{1350}{400}}$ ($= \frac{3}{2}$) or $\sqrt[3]{\frac{400}{1350}}$ ($= \frac{2}{3}$) or $\sqrt[3]{1350} : \sqrt[3]{400}$ or 3:2 (oe)		3	M1 Obtain linear scale factor. This mark not awarded unless cube root taken at some stage.
	$360 \times \left(\sqrt[3]{\frac{1350}{400}} \right)^2$ or 360×2.25 or $360 \div \left(\sqrt[3]{\frac{400}{1350}} \right)^2$ or $360 \div \frac{4}{9}$			M1(DEP)
		810		A1
				Total 3 marks

16	$\frac{(2x-1)(x+5)}{3(2x-1)(2x+1)}$ or $\frac{(2x-1)(x+5)}{(2x-1)(6x+3)}$		3	M1 factorise numerator M1 factorise denominator These marks are independent
		$\frac{x+5}{3(2x+1)}$		A1 accept $\frac{x+5}{6x+3}$
				Total 3 marks

17	(a)	$\frac{40}{32}$ or 75 or $\frac{5}{4}$ or $1\frac{1}{4}$ or 1.25		2	M1
			1 hour 15 minutes		A1
	(b)	$\frac{1.5+40+10}{22.5+75+37.5} \times 60 = \frac{51.5}{135}$ or $\frac{1.5+40+10}{0.375+1.25+0.625} = \frac{51.5}{2.25}$ (oe)		2	M1 FT 75 or 1.25 from (a)
			22.9		A1 awrt 22.9
Total 4 marks					

18	$3x + y = 7$ $3x + 21y = -39$	$21x + 7y = 49$ $x + 7y = -13$	$y = 7 - 3x$	$x = -13 - 7y$	4	M1 Two equations with equal x or y coefficients, or correctly isolate x or y .	Allow a single mistake across both M marks.
	$20y = -46$	$20x = 62$	$x + 7(7 - 3x) = -13$	$3 \times (-13 - 7y) + y = 7$		M1 DEP Correct choice to add or subtract equations, or substitute x or y into other equation	
				$x = 3.1$ (oe)		A1	
				$y = -2.3$ (oe)		A1	
Total 4 marks							

19	$\sin 42 = \frac{5.2}{AC}$		4	M1	Any fully correct equation containing AC
	$AC = \frac{5.2}{\sin 42}$ (= 7.77 or better)			M1 DEP	Any fully correct expression for AC
	$\tan CAD = \frac{4.5}{"7.77"}$			M1 INDEP	ft $AC > 0$ Any fully correct expression for CAD or a trig ratio of CAD.
		30.1		A1	
					Total 4 marks


20	(a)		$2x - \frac{16}{x^2}$	2	B2	B1 for $2x$, B1 for $-\frac{16}{x^2}$ or $-16x^{-2}$
	(b)	$2x - \frac{16}{x^2} = 0$ (oe)		2	M1	$\frac{dy}{dx} = 0$, ft their $\frac{dy}{dx}$ as long as this is not the same as y
			2		A1	CSO
					Total 4 marks	

21	$6 + 2x = 14$		4	M1	
		$x = 4$		A1	
	$3y - 3x = -27$ or $3y - 3 \times "4" = -27$			M1	
		$y = -5$		A1	
	NB award both M marks if $\begin{pmatrix} 6 + 2x \\ 3y - 3x \end{pmatrix} = \begin{pmatrix} 14 \\ -27 \end{pmatrix}$ seen.				Total 4 marks

22	$\text{Ext}_{12} = \frac{360}{12} (=30)$	$\text{Int}_{12} = \frac{(2 \times 12 - 4)90}{12}$ (= 150)		4	M1
	$\text{Ext}_n = 54 - "30" (= 24)$	$\text{Int}_n = 360 - 54 - "150"$ (= 156)			M1 INDEP
	$n = \frac{360}{"24"}$	$\frac{(2n - 4)90}{n} = "156" \text{ (oe)}$			M1 DEP Dependent on both previous method marks
			15		A1
					Total 4 marks

23	(a)		135	1	B1
	(b)	$r^2 - 9 = 46(r + 3)$		3	M1 Allow different variable used.
		$r - 3 - 46 = 0$ or $r - 3 = 46$ or $(r - 49)(r + 3) = 0$ or $(r =) \frac{-(-46) + \sqrt{(-46)^2 - 4 \times 1 \times -147}}{2 \times 1}$			M1 (DEP)
			49		A1 49 and -3 together gain M1M1A0
		NB for factorisation in (b) allow any expression which gives 2 out of 3 correct in $r^2 - 46r - 147$ eg. $(r - 50)(r + 4)$			Total 4 marks

24	(a)	$0.2 + x + x + 0.1 + 0.18 + x + 0.1 = 1,$ $3x + 0.58 = 1 \text{ (oe)}$		2	M1
			0.14 (oe)		A1 $\frac{7}{50}$
	(b)		2	1	B1 ft for $x > 0$
	(c)	60×0.2		2	M1
			12		A1
					Total 5 marks

27	(a) (i)	equal intersecting arcs, centres A and C		4	M1
			perpendicular bisector drawn		A1 Any length.
	(ii)	arc centre A to intersect AC and AB , and equal intersecting arcs from these points of intersection			M1
			angle bisector drawn		A1 Must extend from A to BC .
	(b)	Arc, centre B , radius 5 cm		2	M1
			region R shaded		A1 FT their perpendicular bisector and angle bisector. The area must be bounded by at least both of these and the correct arc.
					Total 6 marks

28	(a) (i)	$2\pi r = \frac{216}{360} \times 2\pi \times 5x$ (oe)		4	M1
			$3x$		A1
	(ii)	$(5x)^2 = ("3x")^2 + h^2$ (oe)			M1
			$4x$		A1
	(b)	$\frac{1}{3}\pi("3x")^2 \times "4x" = \frac{375\pi}{2}$		3	M1 Condone missing brackets for this mark only
		$x^3 = \frac{375 \times 3}{2 \times 9 \times 4} \left(= \frac{125}{8} \right)$ (oe)			M1 DEP
			$2\frac{1}{2}$ (oe)		A1 $\frac{5}{2}, 2.5$
					Total 7 marks

Pearson Education Limited. Registered company number 872828
with its registered office at 80 Strand, London, WC2R 0RL, United Kingdom